

# Conditionally monotone independence

Takahiro Hasebe\*

Graduate School of Science, Kyoto University,

Kyoto 606-8502, Japan

E-mail: hsb@kurims.kyoto-u.ac.jp

## Abstract

We define the notion of conditionally monotone product as a part of conditionally free product, which naturally includes monotone and Boolean products, and moreover, orthogonal product. Then we define conditionally monotone cumulants which are useful to calculate the limit distributions in central limit theorem and Poisson's law of small numbers. We also prove a combinatorial moment-cumulant formula which is parallel to the one for conditionally free convolution. Moreover, we introduce deformed convolutions arising from the conditionally monotone convolution of probability measures and compute the limit distributions. In order to understand the validity of cumulants, we discuss what are cumulants of a given convolution product in general.

## 1 Introduction

Non-commutative notions of independence are connected deeply with the products of states on the free product of algebras (with or without the identification of units). There are two “universal products” (classical and free) defined on the free product of algebras with the identification of units [3, 31] and there are five “natural products” (classical, free, Boolean, monotone and anti-monotone) defined on the free product of algebras without the identification of units [25, 26]. These results can also be understood in terms of tensor structures with inclusions [2].

The conditionally free product of states and the conditionally free convolution of probability measures have been introduced as generalizations of the free product and the free convolution [6]. This notion can be seen as a universal product of pairs of states which could be defined similarly to the definition in [31].

Free and Boolean products and their cumulants [32, 33] can be derived from conditionally free product and conditionally free cumulants. In addition, monotone product can also be derived from conditionally free product [13]. This is an interesting fact, which is, however, not easy to manipulate in the following points: it is difficult to repeat the calculation of monotone product in terms of conditionally free product; it is difficult to derive monotone cumulants [11] from conditionally free cumulants.

In this paper, we introduce conditionally monotone product of pairs of states which is a parallel notion to conditionally free product although it can be defined as a part of conditionally free product. In terms of conditionally monotone product, monotone and Boolean products appear naturally. We also define the notion of conditionally monotone independence as an abstraction of the conditionally monotone product. We briefly mention a connection between conditionally monotone (free) independence and orthogonal independence, the latter of which has been introduced in [19]. As a result, the characterization of orthogonal convolution (Theorem 6.2 in [19]) can be derived from the one of conditionally monotone convolution.

In terms of probability measures, we can define conditionally monotone (additive) convolution. We introduce conditionally monotone cumulants which linearize the powers of a probability measure. Monotone and Boolean cumulants can be derived from conditionally monotone cumulants. We obtain the moment-cumulant formula with the use of the combinatorics of monotone partitions.

---

\*This work was supported by Grant-in-Aid for JSPS Fellows.

Multiplicative conditionally monotone convolution will also be interesting, which is expected to include monotone, Boolean and orthogonal multiplicative convolutions (see [4, 5, 13, 14, 20]). This aspect will be studied in another paper.

In connection to conditionally free convolution, the  $t$ -transformation has been introduced in [7], which is later generalized in [16]. It is important that the  $t$ -deformation of free convolution gives an interpolation between free and Boolean convolutions [7, 8]. Moreover, several other deformations of convolutions arising from conditionally free convolution have been found in [15, 28, 29]. (In [18] another type of interpolation has been obtained.) Motivated by these works, we define deformed convolutions arising from conditionally monotone convolution associated with maps from the set of probability measures into itself. These kinds of convolutions naturally include monotone and Boolean convolutions. Then we show that the generalized  $t$ -transformation, the  $V_a$ -transformation and a transformation related to a  $\triangleright$ -infinitely divisible distribution give associative convolutions in the present situation. In particular, the  $t$ -transformation gives an interpolation between monotone and Boolean convolutions. Thus, many transformations found in [7, 8, 15, 16, 28, 29] also give associative convolutions also in the present situation.

Moreover, we show necessary and sufficient conditions under which the deformed convolution preserves the set  $\{\mu; \text{supp } \mu \subset [0, \infty)\}$  and the set  $\{\mu; \mu \text{ is symmetric}\}$ . These results generalize the results in Boolean and monotone cases [10].

We introduce cumulants of the deformed convolution by a map  $T$ . Then we calculate limit distributions for some associative convolutions. In order to confirm the validity of the cumulants of the deformed convolution, we consider the uniqueness and the existence of cumulants of a general convolution product in Section 4.

## 2 Preliminaries

### 2.1 Reciprocal Cauchy transform

The Cauchy transform of a probability measure  $\mu$  is defined by

$$G_\mu(z) = \int_{\mathbb{R}} \frac{1}{z - x} d\mu(x), \quad z \in \mathbb{C} \setminus \mathbb{R}. \quad (2.1)$$

The reciprocal Cauchy transform of a probability measure  $\mu$  is defined by

$$H_\mu(z) = \frac{1}{G_\mu(z)}, \quad z \in \mathbb{C} \setminus \mathbb{R}, \quad (2.2)$$

which is an analytic function from the upper halfplane into itself. As a result,  $H_\mu$  has the following form:

$$H_\mu(z) = b + z + \int_{\mathbb{R}} \frac{1 + xz}{x - z} d\eta(x), \quad (2.3)$$

where  $b \in \mathbb{R}$  and  $\eta$  is a positive finite measure. Conversely, any function of the form of the right hand side of (2.3) is a reciprocal Cauchy transform of a probability measure.

### 2.2 Monotone independence

Muraki has defined the notion of monotone independence in [24]. The following definition is slightly different from the original one. Let  $(\mathcal{A}, \varphi)$  be an algebraic probability space and let  $I$  be a linearly ordered set. A family of subalgebras  $\{\mathcal{A}_i\}_{i \in I}$  is said to be monotone independent if the equality

$$\varphi(a_1 a_2 \cdots a_n) = \varphi(a_j) \varphi(a_1 a_2 \cdots a_{j-1} a_{j+1} \cdots a_n) \quad (2.4)$$

holds for  $a_k \in \mathcal{A}_{i_k}$  with  $i_1, i_2, \dots, i_n \in I$ ,  $i_{j-1} < i_j > i_{j+1}$  and  $1 \leq j \leq n$ . When  $j = 1$  (resp.  $j = n$ ), the condition  $i_{j-1} < i_j > i_{j+1}$  becomes  $i_1 > i_2$  (resp.  $i_{n-1} < i_n$ ). He has also defined the monotone convolution  $\mu \triangleright \nu$  of two probability measures  $\mu, \nu$  and proved that monotone convolution is characterized by the formula

$$H_{\mu \triangleright \nu} = H_\mu \circ H_\nu. \quad (2.5)$$

Monotone convolution is non-commutative and associative.

Let  $\mathcal{NC}(n)$  be the set of all non-crossing partitions [27]. Let  $\mathcal{M}(n)$  be the set of all *monotone partitions* defined by

$$\mathcal{M}(n) := \{(\pi, \lambda); \pi \in \mathcal{NC}(n), \text{ if } V, W \in \pi \text{ and } V \text{ is in the inner side of } W, \text{ then } V >_\lambda W\}, \quad (2.6)$$

where  $\lambda$  denotes a linear ordering of the blocks of  $\pi$  and  $V >_\lambda W$  means that  $V$  is larger than  $W$  under the linear ordering  $\lambda$ . (See [25, 26], and also [21, 22].)

In the paper [11] the notion of monotone cumulants have been defined. Monotone cumulants do not satisfy the additivity for general probability measures, but satisfy the additivity for identical ones:  $r_n(\mu^{\boxplus N}) = Nr_n(\mu)$ . The moment-cumulant formula is given by

$$m_n(\mu) = \sum_{(\pi, \lambda) \in \mathcal{M}(n)} \frac{1}{|\pi|!} \prod_{V \in \pi} r_{|V|}(\mu). \quad (2.7)$$

### 2.3 Conditionally free independence

Let  $I$  be an index set. Let  $\mathcal{A}_i$  be a unital  $*$ -algebra and let  $\varphi_i, \psi_i$  be states on  $\mathcal{A}_i$  for  $i \in I$ . The conditionally free product of triples  $(\mathcal{A}_i, \varphi_i, \psi_i)_{i \in I}$  has been introduced in the paper [6]. We set  $(\mathcal{A}, \varphi, \psi) = *_{i \in I}(\mathcal{A}_i, \varphi_i, \psi_i)$  by  $\mathcal{A} := *_{i \in I} \mathcal{A}_i$  (the free product with the identification of units) and  $\psi := *_{i \in I} \psi_i$  (the free product of states).  $\varphi$  is defined by the following condition: the equality

$$\varphi(a_1 \cdots a_n) = \prod_{k=1}^n \varphi_{i_k}(a_k) \quad (2.8)$$

holds if  $a_k \in \mathcal{A}_{i_k}$  with  $i_1 \neq \cdots \neq i_n$  and  $\psi_{i_k}(a_k) = 0$  for all  $1 \leq k \leq n$ .

The conditionally free convolution of pairs of probability measures  $(\mu_1, \nu_1)$  and  $(\mu_2, \nu_2)$  also has been introduced in the paper [6]. Define the  $R$ -transform and the conditionally free  $R$ -transform by

$$\frac{1}{G_{\nu_i}(z)} = z - R_{\nu_i}(G_{\nu_i}(z)), \quad (2.9)$$

$$\frac{1}{G_{\mu_i}(z)} = z - R_{(\mu_i, \nu_i)}(G_{\nu_i}(z)), \quad (2.10)$$

for  $i = 1, 2$ . We expand  $R_{(\mu, \nu)}(z) = \sum_{n=1}^{\infty} R_n(\mu, \nu) z^{n-1}$  as a formal power series.  $R_n(\mu, \nu)$  are called conditionally free cumulants. Similarly, we expand  $R_\nu(z) = \sum_{n=1}^{\infty} R_n(\nu) z^{n-1}$  and  $R_n(\nu)$  are called free cumulants. In this paper,  $H_\mu(z)$  is more useful than  $G_\mu(z)$  and correspondingly we use  $\phi_{(\mu, \nu)}(z) := R_{(\mu, \nu)}(\frac{1}{z})$  and  $\phi_\nu(z) := R_\nu(\frac{1}{z})$ . Then (2.9) and (2.10) can be written as

$$H_{\nu_i}(z) = z - \phi_{\nu_i}(H_{\nu_i}(z)), \quad (2.11)$$

$$H_{\mu_i}(z) = z - \phi_{(\mu_i, \nu_i)}(H_{\nu_i}(z)). \quad (2.12)$$

Moment-cumulant formula for conditionally free convolution is given by

$$m_n(\mu) = \sum_{\pi \in \mathcal{NC}(n)} \left( \prod_{\substack{V \in \pi, \\ V: \text{ outer}}} R_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \pi, \\ V: \text{ inner}}} R_{|V|}(\nu) \right), \quad (2.13)$$

which is a generalization of the moment-cumulant formulae for free and Boolean cases.

The conditionally free convolution of  $(\mu_1, \nu_1)$  and  $(\mu_2, \nu_2)$  is the pair  $(\mu, \nu) = (\mu_1, \nu_1) \boxplus (\mu_2, \nu_2)$ , where  $\mu$  and  $\nu$  are characterized by

$$\phi_\nu(z) = \phi_{\nu_1}(z) + \phi_{\nu_2}(z), \quad (2.14)$$

$$\phi_{(\mu, \nu)}(z) = \phi_{(\mu_1, \nu_1)}(z) + \phi_{(\mu_2, \nu_2)}(z). \quad (2.15)$$

We denote by  $(\mu_1 \nu_1 \boxplus \mu_2, \nu_1 \boxplus \nu_2)$  the conditionally free convolution of  $(\mu_1, \nu_1)$  and  $(\mu_2, \nu_2)$ .

## 2.4 Technical facts

We summarize the notation and several lemmata which will be used in this paper. Let  $\mathcal{P}$ ,  $\mathcal{P}^2$ ,  $\mathcal{P}_m$ ,  $\mathcal{P}_c$ ,  $\mathcal{P}_+$  and  $\mathcal{P}_{\text{sym}}$  be the set of probability measures, the set of probability measures with finite variance, the set of probability measures with finite moments of all orders, the set of probability measures with compact support, the set of probability measures whose supports are contained in  $[0, \infty)$  and the set of symmetric probability measures, respectively. The following lemma is a slight generalization of Proposition 2.2 in [23].

**Lemma 2.1.** *A probability measure  $\mu$  belongs to  $\mathcal{P}^2$  if and only if  $H_\mu$  has the representation*

$$H_\mu(z) = a + z + \int_{\mathbb{R}} \frac{1}{x - z} d\rho(x), \quad (2.16)$$

where  $a \in \mathbb{R}$ ,  $\rho$  a positive finite measure.  $a$  and  $\rho$  are determined uniquely. Furthermore, we have  $\rho(\mathbb{R}) = \sigma^2(\mu)$  and  $a = -m(\mu)$ , where  $m(\mu)$  denotes the mean of  $\mu$  and  $\sigma^2(\mu)$  denotes the variance of  $\mu$ .

We define  $a(\mu) := \inf\{x \in \text{supp}\mu\}$  and  $b(\mu) := \sup\{x \in \text{supp}\mu\}$ . We note that  $-\infty \leq a(\mu) < \infty$  and  $-\infty < b(\mu) \leq \infty$  hold. The following lemmata 2.2-2.5 have been proved in [10].

**Lemma 2.2.** *Let  $\nu$  and  $\mu$  be probability measures. Then the following inequalities hold:*

- (1) *If  $\text{supp}\nu \cap (-\infty, 0] \neq \emptyset$  and  $\text{supp}\nu \cap [0, \infty) \neq \emptyset$ , then  $a(\mu) \geq a(\nu \triangleright \mu)$  and  $b(\mu) \leq b(\nu \triangleright \mu)$ .*
- (2) *If  $\text{supp}\nu \subset (-\infty, 0]$ , then  $a(\mu) \geq a(\nu \triangleright \mu)$  and  $b(\nu) + b(\mu) \leq b(\nu \triangleright \mu)$ .*
- (3) *If  $\text{supp}\nu \subset [0, \infty)$ , then  $a(\nu) + a(\mu) \geq a(\nu \triangleright \mu)$  and  $b(\mu) \leq b(\nu \triangleright \mu)$ .*

**Lemma 2.3.** *We use the notation in (2.3). For  $\mu \in \mathcal{P}$ , the condition  $\mu \in \mathcal{P}_+$  is equivalent to the following conditions.*

- (1)  *$\text{supp}\eta \subset [0, \infty)$ ;*
- (2)  *$H_\mu(-0) \leq 0$ .*

*The conditions (2) is equivalent to the conditions  $\eta(\{0\}) = 0$ ,  $\int_0^\infty \frac{1}{x} d\eta(x) < \infty$  and  $b + \int_0^\infty \frac{1}{x} d\eta(x) \leq 0$ .*

**Lemma 2.4.** *Let  $\{\mu_t\}_{t \geq 0}$  be a weakly continuous  $\triangleright$ -convolution semigroup with  $\mu_0 = \delta_0$ . Then the following statements are equivalent:*

- (1) *there exists  $t_0 > 0$  such that  $\text{supp}\mu_{t_0} \subset [0, \infty)$ ;*
- (2)  *$\text{supp}\mu_t \subset [0, \infty)$  for all  $0 \leq t < \infty$ ;*

**Lemma 2.5.** *We assume that the support of each  $\mu_t$  is compact (this is a time-independent property). Then the following statements are equivalent.*

- (1) *There exists  $t_0 > 0$  such that  $\mu_{t_0}$  is symmetric.*
- (2)  *$\mu_t$  is symmetric for all  $t > 0$ .*

**Lemma 2.6.**  *$\mathcal{P}_+$  and  $\mathcal{P}_{\text{sym}}$  are closed subsets of  $\mathcal{P}$  under the weak topology.*

*Proof.* Let  $\{\mu_n\} \subset \mathcal{P}_+$  be a sequence converging to  $\mu \in \mathcal{P}$ . The weak convergence implies that  $\mu((-\infty, 0)) \leq \liminf \mu_n((-\infty, 0)) = 0$ ; therefore,  $\mu \in \mathcal{P}_+$ .

For a probability measure  $\nu$ ,  $\nu \in \mathcal{P}_{\text{sym}}$  is equivalent to the condition

$$\int_{\mathbb{R}} g(x) d\nu(x) = 0 \text{ for all } g \in C_b(\mathbb{R}), \ g(-x) = -g(x), \quad (2.17)$$

where  $C_b(\mathbb{R})$  is the set of bounded continuous functions on  $\mathbb{R}$ . This equivalence can be proved by a simple approximation argument. Then the conclusion is not difficult.  $\square$

### 3 Conditionally monotone independence

It is known that free, Boolean and monotone product of states (denoted by  $*$ ,  $\diamond$  and  $\triangleright$ , respectively) can be derived from conditionally free product [6, 13], which we explain briefly. We consider triples of algebras and states  $(\mathcal{A}_1, \varphi_1, \psi_1)$  and  $(\mathcal{A}_2, \varphi_2, \psi_2)$ . We assume that  $\mathcal{A}_i$  has a decomposition

$$\mathcal{A}_i = \mathbb{C} 1 \oplus \mathcal{A}_i^0 \quad (3.1)$$

with a subalgebra  $\mathcal{A}_i^0$  ( $i = 1, 2$ ). Then the delta state  $\delta_i$  ( $i = 1, 2$ ) is defined by  $\delta_i(\lambda 1 + a^0) = \lambda$  for  $\lambda \in \mathbb{C}$  and  $a^0 \in \mathcal{A}_i^0$ .  $\delta_i$  is a homomorphism from  $\mathcal{A}_i$  to  $\mathbb{C}$ . Conversely, if there exists a homomorphism from  $\mathcal{A}_i$  to  $\mathbb{C}$ , then  $\mathcal{A}_i^0$  is given by its kernel. Then we have the following relations.

$$(\varphi, \varphi) * (\psi, \psi) = (\varphi * \psi, \varphi * \psi) \text{ on } \mathcal{A}_1 * \mathcal{A}_2, \quad (3.2)$$

$$(\varphi, \delta_1) * (\psi, \delta_2) = (\varphi \diamond \psi, \delta_1 * \delta_2) \text{ on } \mathcal{A}_1^0 * \mathcal{A}_2^0, \quad (3.3)$$

$$(\varphi, \delta_1) * (\psi, \psi) = (\varphi \triangleright \psi, \psi) \text{ on } \mathcal{A}_1^0 * \mathcal{A}_2. \quad (3.4)$$

In terms of (compactly supported) probability measures, the equalities (3.2)-(3.4) can be written as

$$(\mu, \mu) \boxplus (\nu, \nu) = (\mu \boxplus \nu, \mu \boxplus \nu), \quad (3.5)$$

$$(\mu, \delta_0) \boxplus (\nu, \delta_0) = (\mu \boxplus \nu, \delta_0), \quad (3.6)$$

$$(\mu, \delta_0) \boxplus (\nu, \nu) = (\mu \triangleright \nu, \nu). \quad (3.7)$$

We can understand the associativity of Boolean convolution by (3.6) since conditionally free convolution is associative. The associativity of monotone convolution is, however, not easy to understand by (3.7) since we cannot repeat the calculation of (3.7). We show that the monotone convolution of the second components of pairs of probability measures is suitable to repeat the calculation.

**Definition 3.1.** (1) Let  $(\mathcal{A}_1, \varphi_1, \psi_1)$  and  $(\mathcal{A}_2, \varphi_2, \psi_2)$  be triples consisting of algebras and two linear functionals and let  $\mathcal{A}_1 *_{nu} \mathcal{A}_2$  be the free product without the identification of units. We define conditionally monotone product

$$(\mathcal{A}_1, \varphi_1, \psi_1) \triangleright (\mathcal{A}_2, \varphi_2, \psi_2) := (\mathcal{A}_1 *_{nu} \mathcal{A}_2, \varphi_1 \triangleright_{\psi_2} \varphi_2, \psi_1 \triangleright \psi_2), \quad (3.8)$$

where  $\varphi_1 \triangleright_{\psi_2} \varphi_2$  is defined as follows. We denote the unitization of each  $\mathcal{A}_i$  by  $\widetilde{\mathcal{A}}_i := \mathbb{C} 1_{\widetilde{\mathcal{A}}_i} \oplus \mathcal{A}_i$  (see [2]). (When  $\mathcal{A}_i$  contains a unit  $1_{\mathcal{A}_i}$ ,  $1_{\mathcal{A}_i} \neq 1_{\widetilde{\mathcal{A}}_i}$ .) Then we have  $\widetilde{\mathcal{A}}_1 *_{nu} \widetilde{\mathcal{A}}_2 \cong \widetilde{\mathcal{A}}_1 * \widetilde{\mathcal{A}}_2$  naturally. The linear functionals  $\varphi_i$  and  $\psi_i$  are naturally extended to  $\widetilde{\varphi}_i$  and  $\widetilde{\psi}_i$  on  $\widetilde{\mathcal{A}}_i$  by  $\widetilde{\psi}_i(1_{\widetilde{\mathcal{A}}_i}) = 1$  and  $\widetilde{\psi}_i(1_{\mathcal{A}_i}) = 1$ . We denote by  $\widetilde{\delta}_1$  the delta state of the decomposition  $\widetilde{\mathcal{A}}_1 = \mathbb{C} 1 \oplus \mathcal{A}_1$ . We define  $\varphi_1 \triangleright_{\psi_2} \varphi_2$  by the restriction of  $\widetilde{\varphi}_1 \widetilde{\delta}_1 *_{\widetilde{\psi}_2} \widetilde{\varphi}_2$  on  $\mathcal{A}_1 *_{nu} \mathcal{A}_2$ .

(2) For pairs of probability measures  $(\mu_1, \nu_1)$  and  $(\mu_2, \nu_2)$ , we define the conditionally monotone convolution  $(\mu_1, \nu_1) \triangleright (\mu_2, \nu_2)$  by  $(\mu_1, \nu_1) \triangleright (\mu_2, \nu_2) := (\mu_1 \triangleright_{\nu_2} \mu_2, \nu_1 \triangleright \nu_2)$ , where  $\mu_1 \triangleright_{\nu_2} \mu_2$  is defined by  $\mu_1 \triangleright_{\nu_2} \mu_2 := \mu_1 \delta_0 \boxplus_{\nu_2} \mu_2$ .

**Remark 3.2.** Since conditionally monotone product is defined on the free product without the identification of units, the definition has been given for linear functionals. Conditionally monotone product, however, preserves the positivity of linear functionals since it is defined by the restriction of conditionally free product.

**Proposition 3.3.**  $\mu_1 \triangleright_{\nu_2} \mu_2$  is characterized by

$$H_{\mu_1 \triangleright_{\nu_2} \mu_2} = H_{\mu_1} \circ H_{\nu_2} + H_{\mu_2} - H_{\nu_2}. \quad (3.9)$$

*Proof.* We immediately obtain the equality  $H_{\mu_1} \circ H_{\nu_2}(z) = H_{\nu_2}(z) - \phi_{(\mu_1, \delta_0)}(H_{\nu_2}(z))$  by (2.12). Since conditionally free product is characterized by the sum of  $\phi_{(\cdot)}$ , we have  $H_{\mu_1} \circ H_{\nu_2}(z) - H_{\nu_2}(z) + H_{\mu_2}(z) = z - \phi_{(\mu_1 \triangleright_{\nu_2} \mu_2, \nu_2)} \circ H_{\nu_2}(z)$ .  $\square$

We can easily check by Proposition 3.3 that the conditionally monotone convolution of probability measures is associative, i.e.,  $((\mu_1, \nu_1) \triangleright (\mu_2, \nu_2)) \triangleright (\mu_3, \nu_3) = (\mu_1, \nu_1) \triangleright ((\mu_2, \nu_2) \triangleright (\mu_3, \nu_3))$ . Then it is expected to be the case that the conditionally monotone product of pairs of linear functionals is also associative, which will be proved below. We use the following notation: for a linearly ordered index set  $I = \{i_1, i_2, \dots, i_n\}$  with  $i_1 < i_2 < \dots < i_n$ , we set

$$\overrightarrow{\prod_{i \in I} a_i} := a_{i_1} a_{i_2} \cdots a_{i_n}.$$

**Proposition 3.4.** *Let  $(\mathcal{A}_1, \varphi_1, \psi_1)$  and  $(\mathcal{A}_2, \varphi_2, \psi_2)$  be triples consisting of algebras and two linear functionals. We have the following formula.*

$$\varphi_1 \triangleright_{\psi_2} \varphi_2(a_1 b_1 a_2 \cdots b_{n-1} a_n) = \sum_{S \subset \{1, \dots, n-1\}} \left( \prod_{j \notin S} \psi_2(b_j) \right) \left( \prod_{j \in S} (\varphi_2(b_j) - \psi_2(b_j)) \right) \prod_j \varphi_1(a_{S_j}) \quad (3.10)$$

for  $a_k \in \mathcal{A}_1$ ,  $b_k \in \mathcal{A}_2$ , where  $a_{S_j} := \overrightarrow{\prod_{k \in S_j} a_k}$ .  $\{S_j\}$  is a partition of  $\{1, \dots, n\}$  determined by  $S$  as follows: if  $S = \{k_1, \dots, k_m\}$  with  $1 \leq k_1 < \dots < k_m \leq n-1$ , then  $S_j := \{k_{j-1} + 1, \dots, k_j\}$  for  $1 \leq j \leq m+1$ , where  $k_0 := 0$  and  $k_{m+1} := n$ . If  $S = \emptyset$ , then  $S_1 = \{1, \dots, n\}$ .  $\prod_{j \in \emptyset} \psi_2(b_j)$  is defined to be 1.

*Proof.* Let  $x_j$  and  $y_k$  be (non-commutative) variables in an algebra over  $\mathbb{C}$  with unit 1 and let  $p_j \in \mathbb{C}$ . Then the following identity holds:

$$x_1 y_1 x_2 \cdots y_{n-1} x_n = \sum_{S \subset \{1, \dots, n-1\}} \left( \prod_{j \notin S} p_j \right) \left( x_{S_1} (y_{k_1} - p_{k_1} 1) x_{S_2} (y_{k_2} - p_{k_2} 1) \cdots (y_{k_m} - p_{k_m} 1) x_{S_{m+1}} \right) \quad (3.11)$$

where we have used the same notation as explained below (3.10). (3.11) is proved by induction about the number  $n$  easily. We put  $x_j = a_j \in \mathcal{A}_1$ ,  $y_j = b_j \in \mathcal{A}_2$ ,  $p_j = \psi_2(b_j)$  and we use the unit  $1_{\widetilde{\mathcal{A}_1 *_{nu} \mathcal{A}_2}}$ . Taking the expectation of (3.11) w.r.t.  $\widetilde{\varphi_1}_{\widetilde{\delta_1}} *_{\widetilde{\psi_2}} \widetilde{\varphi_2}$ , we obtain the result by the definition of the conditionally free product.  $\square$

**Theorem 3.5.** *Let  $(\mathcal{A}_1, \varphi_1, \psi_1)$  and  $(\mathcal{A}_2, \varphi_2, \psi_2)$  be triples consisting of algebras and two linear functionals. The calculation rules for the conditionally monotone product are what follows.*

- (1)  $\varphi_1 \triangleright_{\psi_2} \varphi_2(b_1 a_1 b_2 a_2 \cdots a_n b_{n+1}) = \varphi_2(b_1) \varphi_2(b_{n+1}) \varphi_1 \triangleright_{\psi_2} \varphi_2(a_1 b_2 a_2 \cdots b_n a_n)$   
for  $a_k \in \mathcal{A}_1$ ,  $b_k \in \mathcal{A}_2$ ,  $n \in \mathbb{N}$ .
- (2)  $\varphi_1 \triangleright_{\psi_2} \varphi_2(a_1 b_1 \cdots b_{n-1} a_n) = (\varphi_2(b_j) - \psi_2(b_j)) \varphi_1 \triangleright_{\psi_2} \varphi_2(a_1 b_1 a_2 \cdots b_{j-1} a_j) \varphi_1 \triangleright_{\psi_2} \varphi_2(a_{j+1} b_{j+1} \cdots b_{n-1} a_n)$   
+  $\psi_2(b_j) \varphi_1 \triangleright_{\psi_2} \varphi_2(a_1 b_1 a_2 \cdots b_{j-1} a_j a_{j+1} b_{j+1} \cdots b_{n-1} a_n)$   
for  $a_k \in \mathcal{A}_1$ ,  $b_k \in \mathcal{A}_2$ ,  $1 \leq j \leq n-1$ ,  $n \in \mathbb{N}$ .

*Proof.* Since (1) can be proved similarly, we only prove (2). Moreover, we assume that  $j = 1$  since the proof for general  $j$  is essentially the same. We denote by 1 the unit  $1_{\widetilde{\mathcal{A}_1 *_{nu} \mathcal{A}_2}}$  for simplicity. First we obtain

$$\begin{aligned} \widetilde{\varphi_1}_{\widetilde{\delta_1}} *_{\widetilde{\psi_2}} \widetilde{\varphi_2}(a_1 b_1 a_2 \cdots b_{n-1} a_n) &= \widetilde{\varphi_1}_{\widetilde{\delta_1}} *_{\widetilde{\psi_2}} \widetilde{\varphi_2}(a_1 (b_1 - \psi_2(b_1) 1) a_2 \cdots b_{n-1} a_n) \\ &\quad + \psi_2(b_1) \widetilde{\varphi_1}_{\widetilde{\delta_1}} *_{\widetilde{\psi_2}} \widetilde{\varphi_2}(a_1 a_2 \cdots b_{n-1} a_n). \end{aligned} \quad (3.12)$$

Using the identity (3.11) and the definition of conditionally free product, we have

$$\begin{aligned} &\widetilde{\varphi_1}_{\widetilde{\delta_1}} *_{\widetilde{\psi_2}} \widetilde{\varphi_2}(a_1 (b_1 - \psi_2(b_1) 1) a_2 \cdots b_{n-1} a_n) \\ &= \widetilde{\varphi_1}_{\widetilde{\delta_1}} *_{\widetilde{\psi_2}} \widetilde{\varphi_2} \left( a_1 (b_1 - \psi_2(b_1) 1) \sum_{S \subset \{2, \dots, n-1\}} \left( \prod_{j \notin S} \psi_2(b_j) \right) \left( a_{S_1} (b_{k_1} - \psi_2(b_{k_1}) 1) \cdots a_{S_{m+1}} \right) \right) \\ &= (\varphi_2(b_1) - \psi_2(b_1)) \sum_{S \subset \{2, \dots, n-1\}} \left( \prod_{j \notin S} \psi_2(b_j) \right) \left( \prod_{j \in S} (\varphi_2(b_j) - \psi_2(b_j)) \right) \varphi_1(a_1) \prod_{j=1}^{|S|+1} \varphi_1(a_{S_j}) \\ &= \varphi_1(a_1) (\varphi_2(b_1) - \psi_2(b_1)) \widetilde{\varphi_1}_{\widetilde{\delta_1}} *_{\widetilde{\psi_2}} \widetilde{\varphi_2}(a_2 b_2 \cdots b_{n-1} a_n). \end{aligned}$$

$\square$

**Theorem 3.6.** *Conditionally monotone product is associative.*

*Proof.* Let  $(\mathcal{A}_i, \varphi_i, \psi_i)$  ( $1 \leq i \leq 3$ ) be triples consisting of algebras and two linear functionals. We identify  $(\mathcal{A}_1 *_{nu} \mathcal{A}_2) *_{nu} \mathcal{A}_3$  with  $\mathcal{A}_1 *_{nu} (\mathcal{A}_2 *_{nu} \mathcal{A}_3)$  naturally and denote them by  $\mathcal{A}_1 *_{nu} \mathcal{A}_2 *_{nu} \mathcal{A}_3$ . What should be proved is the equality  $\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3) = (\varphi_1 \triangleright_{\psi_2} \varphi_2) \triangleright_{\psi_3} \varphi_3$  on  $\mathcal{A}_1 *_{nu} \mathcal{A}_2 *_{nu} \mathcal{A}_3$ . We call  $x \in \mathcal{A}_1 *_{nu} \mathcal{A}_2 *_{nu} \mathcal{A}_3$  an elementary element if  $x$  is of such a form as  $x = x_1 x_2 \cdots x_N$ ,  $x_j \in \mathcal{A}_{i_j}$ ,  $i_1 \neq \cdots \neq i_n$ . We give a proof by induction about the number of elements of  $\mathcal{A}_3$  in elementary elements. In this proof we use the notation  $a, a'$  or  $a_j$  for elements in  $\mathcal{A}_1$ ,  $b, b'$  or  $b_j$  for elements in  $\mathcal{A}_2$  and  $c$  or  $c_j$  for elements in  $\mathcal{A}_3$  without any references.

When  $x \in \mathcal{A}_2 *_{nu} \mathcal{A}_3 \subset \mathcal{A}_1 *_{nu} \mathcal{A}_2 *_{nu} \mathcal{A}_3$ , we can easily prove that  $\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(x) = (\varphi_1 \triangleright_{\psi_2} \varphi_2) \triangleright_{\psi_3} \varphi_3(x)$ . We assume that the statement is the case for elementary elements containing not more than  $n$  elements in  $\mathcal{A}_3$ . Let  $x$  be an elementary element containing  $n+1$  elements in  $\mathcal{A}_3$ . Let  $c$  be the  $(n+1)$ -th element in  $\mathcal{A}_3$  contained in  $x$ . Except for some cases explained below,  $x$  can be written as  $x = uavcw$ , where each component satisfies the following property:

- (i)  $u$  is an elementary element in  $\mathcal{A}_1 *_{nu} \mathcal{A}_2 *_{nu} \mathcal{A}_3$  containing not more than  $n$  elements in  $\mathcal{A}_3$ ;
- (ii)  $v$  is an elementary element in  $\mathcal{A}_2 *_{nu} \mathcal{A}_3$  containing not more than  $n$  elements in  $\mathcal{A}_3$ . The last element in  $v$  belongs to  $\mathcal{A}_2$ , i.e.,  $v = v'b_1$ ;
- (iii)  $w$  is an elementary element in  $\mathcal{A}_1 *_{nu} \mathcal{A}_2$ .

When  $x$  is of such a form as  $x = uacw$ , the following proof is applicable if  $v$  is understood to play the role of a unit. By the property (1) in Theorem 3.5, the proof is easy when  $x$  is of such a form as  $x = uavc$ . When  $a$  does not appear, again by the property (1) in Theorem 3.5 the proof becomes easy. Therefore, we only prove for  $x$  of the form  $uavcw$ . We consider the following two cases: (1) The first element in  $w$  belongs to  $\mathcal{A}_1$ , i.e.,  $w = a'w'$ ; (2) The first element in  $w$  belongs to  $\mathcal{A}_2$ , i.e.,  $w = b_2w'$ .

Case (1): We obtain

$$\begin{aligned}
& (\varphi_1 \triangleright_{\psi_2} \varphi_2) \triangleright_{\psi_3} \varphi_3(uavcw) \\
&= (\varphi_3(c) - \psi_3(c))(\varphi_1 \triangleright_{\psi_2} \varphi_2) \triangleright_{\psi_3} \varphi_3(uav)(\varphi_1 \triangleright_{\psi_2} \varphi_2) \triangleright_{\psi_3} \varphi_3(w) \\
&\quad + \psi_3(c)(\varphi_1 \triangleright_{\psi_2} \varphi_2) \triangleright_{\psi_3} \varphi_3(uavw) \\
&= (\varphi_3(c) - \psi_3(c))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uav)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad + \psi_3(c)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uavw) \\
&= (\varphi_3(c) - \psi_3(c))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uav)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad + \psi_3(c)(\varphi_2 \triangleright_{\psi_3} \varphi_3(v) - \psi_2 \triangleright \psi_3(v))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad + \psi_3(c)\psi_2 \triangleright \psi_3(v)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uaw),
\end{aligned} \tag{3.13}$$

where we have used the assumption of induction and the property (2) in Theorem 3.5,. On the other hand, we obtain

$$\begin{aligned}
& \varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uavcw) \\
&= (\varphi_2 \triangleright_{\psi_3} \varphi_3(vc) - \psi_2 \triangleright \psi_3(vc))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(w) \\
&\quad + \psi_2 \triangleright \psi_3(vc)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uaw) \\
&= \varphi_2 \triangleright_{\psi_3} \varphi_3(v)\varphi_3(c)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(w) \\
&\quad - \psi_2 \triangleright \psi_3(v)\psi_3(c)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(w) \\
&\quad + \psi_3(c)\psi_2 \triangleright \psi_3(v)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uaw) \\
&= (\varphi_3(c) - \psi_3(c) + \psi_3(c))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uav)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad - \psi_2 \triangleright \psi_3(v)\psi_3(c)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad + \psi_3(c)\psi_2 \triangleright \psi_3(v)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uaw) \\
&= (\varphi_3(c) - \psi_3(c))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uav)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad + \psi_3(c)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\left(\varphi_2 \triangleright_{\psi_3} \varphi_3(v) - \psi_2 \triangleright \psi_3(v)\right)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad + \psi_3(c)\psi_2 \triangleright \psi_3(v)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uaw).
\end{aligned} \tag{3.14}$$

Therefore, they coincide with each other.

Case (2): By the similar calculation, we have

$$\begin{aligned}
& (\varphi_1 \triangleright_{\psi_2} \varphi_2) \triangleright_{\psi_3} \varphi_3(uavcw) \\
&= (\varphi_3(c) - \psi_3(c))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uav)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad + \psi_3(c)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uavw) \\
&= (\varphi_3(c) - \psi_3(c))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_2 \triangleright_{\psi_3} \varphi_3(v)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad + \psi_3(c)(\varphi_2 \triangleright_{\psi_3} \varphi_3(vb_2) - \psi_2 \triangleright \psi_3(vb_2))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w') \\
&\quad + \psi_3(c)\psi_2 \triangleright \psi_3(vb_2)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uaw').
\end{aligned} \tag{3.15}$$

On the other hand we have

$$\begin{aligned}
& \varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uavcw) \\
&= (\varphi_2 \triangleright_{\psi_3} \varphi_3(vcb_2) - \psi_2 \triangleright \psi_3(vcb_2))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w') \\
&\quad + \psi_2 \triangleright \psi_3(vcb_2)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uaw') \\
&= \varphi_2 \triangleright_{\psi_3} \varphi_3(v)(\varphi_3(c) - \psi_3(c))\varphi_2(b_2)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w') \\
&\quad + \psi_3(c)\varphi_2 \triangleright_{\psi_3} \varphi_3(vb_2)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w') \\
&\quad - \psi_2 \triangleright \psi_3(vcb_2)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w') \\
&\quad + \psi_2 \triangleright \psi_3(vcb_2)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uaw') \\
&= \varphi_2 \triangleright_{\psi_3} \varphi_3(v)(\varphi_3(c) - \psi_3(c))\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w) \\
&\quad + \psi_3(c)\varphi_2 \triangleright_{\psi_3} \varphi_3(vb_2)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w') \\
&\quad - \psi_3(c)\psi_2 \triangleright \psi_3(vb_2)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(ua)\varphi_1 \triangleright_{\psi_2} \varphi_2(w') \\
&\quad + \psi_3(c)\psi_2 \triangleright \psi_3(vb_2)\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(uaw').
\end{aligned} \tag{3.16}$$

Then  $\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(x) = (\varphi_1 \triangleright_{\psi_2} \varphi_2) \triangleright_{\psi_3} \varphi_3(x)$  for  $x$  containing  $n + 1$  elements in  $\mathcal{A}_3$ . By induction,  $\varphi_1 \triangleright_{\psi_2 \triangleright \psi_3} (\varphi_2 \triangleright_{\psi_3} \varphi_3)(x) = (\varphi_1 \triangleright_{\psi_2} \varphi_2) \triangleright_{\psi_3} \varphi_3(x)$  holds for all elementary elements  $x$  in  $\mathcal{A}_1 *_{nu} \mathcal{A}_2 *_{nu} \mathcal{A}_3$ .  $\square$

By the associativity of conditionally monotone product, we can define the product  $\triangleright_{i \in I}(\mathcal{A}_i, \varphi_i, \psi_i)$  for a linearly ordered set  $I$ .

Now we define the notion of conditionally monotone independence.

**Definition 3.7.** Let  $(\mathcal{A}, \varphi, \psi)$  be an algebraic probability space equipped with two states. Now we assume that  $\mathcal{A}$  is unital. Let  $I$  be a linearly ordered set. A family of subalgebras  $\{\mathcal{A}_i\}_{i \in I}$  is said to be conditionally monotone independent if the following properties are satisfied:

- (1)  $\varphi(a_1 a_2 \cdots a_n) = \varphi(a_1)\varphi(a_2 \cdots a_n)$  for  $a_k \in \mathcal{A}_{i_k}$  with  $i_1, i_2, \dots, i_n \in I$  and  $i_1 > i_2$ ;
- (2)  $\varphi(a_1 a_2 \cdots a_n) = \varphi(a_n)\varphi(a_1 \cdots a_{n-1})$  for  $a_k \in \mathcal{A}_{i_k}$  with  $i_1, i_2, \dots, i_n \in I$  and  $i_n > i_{n-1}$ ;
- (3)  $\varphi(a_1 a_2 \cdots a_n) = (\varphi(a_j) - \psi(a_j))\varphi(a_1 \cdots a_{j-1})\varphi(a_{j+1} \cdots a_n) + \psi(a_j)\varphi(a_1 a_2 \cdots a_{j-1} a_{j+1} \cdots a_n)$  for  $a_k \in \mathcal{A}_{i_k}$  with  $i_1, i_2, \dots, i_n \in I$  and  $i_{j-1} < i_j > i_{j+1}$  with  $2 \leq j \leq n-1$ .
- (4)  $\psi(a_1 a_2 \cdots a_n) = \psi(a_j)\psi(a_1 a_2 \cdots a_{j-1} a_{j+1} \cdots a_n)$  holds for  $a_k \in \mathcal{A}_{i_k}$  with  $i_1, i_2, \dots, i_n \in I$ ,  $i_{j-1} < i_j > i_{j+1}$  and  $1 \leq j \leq n$ . When  $j = 1$  (resp.  $j = n$ ), the condition  $i_{j-1} < i_j > i_{j+1}$  is understood to be  $i_1 > i_2$  (resp.  $i_{n-1} < i_n$ ).

**Remark 3.8.** This definition includes monotone (resp. Boolean) independence when  $\varphi = \psi$  (resp.  $\psi = 0$ ) on  $\mathcal{A}_i$  for all  $i \in I$ .

We can prove that  $\mathcal{A}_i$  ( $i \in I$ ) are conditionally monotone independent in  $(\mathcal{A}, \varphi, \psi) = \triangleright_{i \in I}(\mathcal{A}_i, \varphi_i, \psi_i)$ .

Here we mention a connection between conditionally monotone/free independence and orthogonal independence. The reader is referred to [19] for the definition and the properties of orthogonal independence and orthogonal (additive) convolution.



**Proposition 3.9.** *In the notation of Definition 3.1,  $\mathcal{A}_1$  is orthogonal to  $\mathcal{A}_2$  with respect to the linear functional  $(\varphi_1 \triangleright_{\psi_2} 0_2, \psi_1 \triangleright \psi_2)$  in the algebra  $\mathcal{A} := \mathcal{A}_1 *_{nu} \mathcal{A}_2$  for arbitrary linear functionals  $\varphi_1, \psi_1, \psi_2$ .  $0_2$  is defined by the 0 linear functional on  $\mathcal{A}_2$ .*

**Remark 3.10.** Orthogonal independence has been defined in a unital algebra in [19]. To state this proposition strictly along the line of the original definition, we only need to unitize  $\mathcal{A}$  and extend the linear functionals to  $\tilde{\mathcal{A}}$ .

*Proof.* This fact follows from Theorem 3.5 immediately.  $\square$

In terms of convolutions of probability measures, we obtain the following relations.

$$(\mu, \delta_0) \boxplus (\delta_0, \nu) = (\mu \vdash \nu, \nu), \quad (3.17)$$

$$(\mu, \lambda) \triangleright (\delta_0, \nu) = (\mu \vdash \nu, \lambda \triangleright \nu). \quad (3.18)$$

Orthogonal convolution has been characterized by  $H_{\mu \vdash \nu}(z) = H_\mu(H_\nu(z)) - H_\nu(z) + z$ . (See Theorem 6.2 in [19].) If we use the relation in Proposition 3.3, this characterization follows immediately.

We give another application. The equality  $\mu \triangleright \nu = (\mu \vdash \nu) \boxplus \nu$  (see Corollary 6.6 in [19]) can be understood by the associativity of conditionally monotone convolution. We first note that  $(\delta_0, \nu) \triangleright (\nu, \delta_0) = (\delta_0 \boxplus \delta_0, \nu \triangleright \nu) = (\nu, \nu)$  and  $(\mu, \mu) \triangleright (\delta_0, \nu) = (\mu \vdash \nu, \mu)$ . Then by the associativity we can calculate  $(\mu, \mu) \triangleright (\delta_0, \nu) \triangleright (\nu, \delta_0)$  in different two ways and we obtain  $((\mu \vdash \nu) \boxplus \nu, \mu \triangleright \nu) = (\mu \triangleright \nu, \mu \triangleright \nu)$ .

## 4 Cumulants of a convolution

To define conditionally monotone cumulants and the related cumulants in the sections 5 and 10, we consider what are cumulants of a convolution product. We do not consider cumulants of a convolution of pairs of probability measures, which can be also possible with a slight generalization.

For a given convolution  $\square$  defined on  $\mathcal{P}_m$ , we consider what are cumulants of  $\square$ . We shall treat convolutions which are not necessarily commutative for the later applications. All results in this section hold for both  $\mathcal{P}_m$  and  $\mathcal{P}_c$ , except for Theorem 4.9. Then we use the set  $\mathcal{P}_m$  mainly.

**Definition 4.1.** (1) We define recursively  $\nu^{\square n} := \nu \square \nu^{\square n-1}$  for  $\nu \in \mathcal{P}_m$ .  $\square$  is said to be power associative if  $\nu^{\square(n+m)} = \nu^{\square n} \square \nu^{\square m}$  for all  $m, n \geq 0$ .

Let  $m_n(\mu)$  be the  $n$ -th moment of  $\mu \in \mathcal{P}_m$ . We put the following assumptions.

(M1) There exists a polynomial  $P_n$  of  $2n - 2$  variables for each  $n \geq 1$  such that

$$m_n(\mu \square \nu) = m_n(\mu) + m_n(\nu) + P_n(m_1(\mu), \dots, m_{n-1}(\mu), m_1(\nu), \dots, m_{n-1}(\nu)). \quad (4.1)$$

(M2) The polynomial  $P_n$  contains no constants for any  $n \geq 1$ .

**Remark 4.2.** The condition (M2) is equal to the condition  $\delta_0 \square \delta_0 = \delta_0$ .

Let  $r_n(\mu)$  be a polynomial of  $\{m_k(\mu)\}_{k \geq 1}$  for any  $n \geq 1$ . We consider the following properties.

(C1) Power additivity: for any  $n, N \geq 1$ ,

$$r_n(\mu^{\square N}) = N r_n(\mu). \quad (4.2)$$

(C2) There exists a polynomial  $Q_n$  of  $n - 1$  variables such that

$$r_n(\mu) = m_n(\mu) + Q_n(m_1(\mu), \dots, m_{n-1}(\mu)). \quad (4.3)$$

(C2') In addition to the condition (C2), the polynomial  $Q_n$  never contains linear terms  $m_k(\mu)$ ,  $1 \leq k \leq n - 1$  for any  $n$ .

If a sequence  $\{r_n\}$  satisfies (C2), we can write  $m_n$  by  $r_n$  as

$$m_n(\mu) = r_n(\mu) + R_n(r_1(\mu), \dots, r_{n-1}(\mu)), \quad (4.4)$$

where  $R_n$  is a polynomial of  $n - 1$  variables.

We note that in many important examples the condition of homogeneity

$$r_n(D_\lambda \mu) = \lambda^n r_n(\mu) \quad (4.5)$$

holds, where  $D_\lambda$  is the dilation operator defined by  $(D_\lambda \mu)(B) = \mu(\lambda^{-1} B)$  (see [17]). Indeed, this condition holds for classical, free, Boolean, monotone convolutions (see [17] for Boolean and free convolutions and see [11] for monotone convolution). Clearly (C2) and (4.5) imply (C2'). We do not assume this condition since the uniqueness of cumulants follows from only (C1) and (C2') (see Theorem 4.4). Moreover, there are examples which satisfy (C2') but do not satisfy (4.5) such as the cumulants of the convolution deformed by the  $V_a$ -transformation [15].

If there exists a sequence  $\{r_n\}$  satisfying (C1) and (C2), we consider a transformation of the form

$$r_n \mapsto r'_n := r_n + \sum_{k=1}^{n-1} a_{n,k} r_k \quad (4.6)$$

for real numbers  $a_{n,k}$ ,  $1 \leq k \leq n-1$ ,  $1 \leq n < \infty$ . This transformation clearly preserves the properties (C1) and (C2). Moreover, we obtain the following property (1).

**Proposition 4.3.** *We assume (M1) and (M2) and assume that  $\square$  is power associative.*

(1) *If there are two sequences  $\{r_n\}$  and  $\{r'_n\}$  satisfying (C1) and (C2), there exists a unique transformation of the form (4.6) which maps  $\{r_n\}$  to  $\{r'_n\}$ .*

(2) *The polynomial  $Q_n$  in (C2) never contains a constant term.*

*Proof.* (1) There exists a polynomial  $A_n$  of variables  $n - 1$  for each  $n \geq 1$  such that  $r'_n = r_n + A_n(r_1, \dots, r_{n-1})$  by (4.3) and (4.4). Replacing  $\mu$  by  $\mu^{\square N}$ , we obtain  $Nr'_n = Nr_n + A_n(Nr_1, \dots, Nr_{n-1})$  for any  $N$ . This is an equality of polynomials of  $N$ , and hence,  $A_n$  is of the form  $A_n(r_1, \dots, r_{n-1}) = \sum_{k=1}^{n-1} a_{n,k} r_k$ .

(2) We show the fact inductively. For  $n = 1$ , there exists  $b_1 \in \mathbb{R}$  such that  $r_1 = m_1 + b_1$ . Since  $P_n$  does not contain a constant term in (4.1), we have  $m_1(\mu \square \mu) = 2m_1(\mu)$ , which implies  $2r_1(\mu) - b_1 = 2r_1(\mu) - 2b_1$ . Therefore,  $b_1 = 0$ . We assume that  $Q_n$  does not contain a constant term for  $n \leq k$ . By a similar argument, we can prove that  $Q_{k+1}$  does not contain a constant term.  $\square$

**Proposition 4.4.** *We assume (M1) and (M2) and assume that  $\square$  is power associative. The following statements are equivalent:*

- (a) *There exists a sequence  $\{r_n\}_{n \geq 1}$  satisfying (C1) and (C2);*
- (b) *There exists a sequence  $\{r_n\}_{n \geq 1}$  satisfying (C1) and (C2');*
- (c)  *$m_n(\mu^{\square N})$  is a polynomial of  $m_1(\mu), \dots, m_n(\mu)$  and  $N$  for any  $n$ .*

Moreover, the sequence  $\{r_n\}$  in (b) is unique and is given by

$$r_n(\mu) = \frac{\partial}{\partial N} m_n(\mu^{\square N}) \Big|_{N=0}. \quad (4.7)$$

**Remark 4.5.** We can see from (4.7) that cumulants are deeply connected with convolution semigroups  $\{\mu_t\}_{t \geq 0}$  with  $\mu_0 = \delta_0$  and the notion of infinite divisibility.

*Proof.* (1) (a)  $\Rightarrow$  (c): If there exists a sequence  $\{r_n\}_{n \geq 1}$  satisfying (C1) and (C2), we have

$$\begin{aligned} m_n(\mu^{\square N}) &= r_n(\mu^{\square N}) + R_n(r_1(\mu^{\square N}), \dots, r_{n-1}(\mu^{\square N})) \\ &= Nr_n(\mu) + R_n(Nr_1(\mu), \dots, Nr_{n-1}(\mu)) \\ &= Nm_n(\mu) + NQ_n(m_1(\mu), \dots, m_{n-1}(\mu)) \\ &\quad + R_n(Nm_1(\mu), \dots, Nm_{n-1}(\mu) + NQ_{n-1}(m_1(\mu), \dots, m_{n-2}(\mu))). \end{aligned} \quad (4.8)$$

Therefore,  $m_n(\mu^{\square N})$  is a polynomial of  $N$  and  $m_k(\mu)$ .

(c)  $\Rightarrow$  (a): By (M1), (M2) and the assumption (c),  $m_n(\mu^{\square N})$  has such a form as

$$m_n(\mu^{\square N}) = Nm_n(\mu) + \sum_{l=0}^L N^l S_l(m_1(\mu), \dots, m_{n-1}(\mu)). \quad (4.9)$$

We define

$$\begin{aligned} r_n(\mu) &:= \left. \frac{\partial}{\partial N} m_n(\mu^{\square N}) \right|_{N=0} \\ &= m_n(\mu) + S_1(m_1(\mu), \dots, m_{n-1}(\mu)). \end{aligned} \quad (4.10)$$

By the power associativity of  $\square$  we obtain (C1). (C2) follows from (4.10).

(a)  $\Rightarrow$  (b): For a sequence  $\{r_n\}$  satisfying (C1) and (C2), we can write  $r_n$  in the form  $r_n = m_n + \sum_{k=1}^{n-1} b_{n,k} m_k + T_n(m_1, \dots, m_{n-1})$ , where  $T_n$  is a polynomial which does not contain linear terms  $m_k$ ,  $1 \leq k \leq n-1$ . We define a new sequence  $\{r'_n\}$  inductively as follows:  $r'_1 := r_1$ ,  $r'_2 = r_2 - b_{2,1} r_1$ ,  $r'_n = r_n - \sum_{k=1}^{n-1} a_{n,k} r'_k$  for  $n \geq 2$ . Then  $r'_n$  do not contain linear terms  $m_k$ .

We note that  $Q_n$  does not contain a constant term by Proposition 4.3 (2). If there exists a sequence  $\{r_n\}$  satisfying (C1) and (C2'), the corresponding polynomial  $R_n$  in (4.4) also does not contain linear terms  $m_k$ ,  $1 \leq k \leq n-1$  and a constant term. Therefore, the equality  $m_n(\mu^{\square N}) = Nr_n(\mu) + R_n(Nr_1(\mu), \dots, Nr_{n-1}(\mu))$  implies that  $r_n = \left. \frac{\partial}{\partial N} m_n(\mu^{\square N}) \right|_{N=0}$ .  $\square$

**Definition 4.6.** Let  $\square$  be a power associative convolution defined on  $\mathcal{P}_m$  satisfying (M1) and (M2). Then the polynomials  $r_n$  satisfying (C1) and (C2') are called the cumulants of the convolution  $\square$ . Cumulants are unique.

**Remark 4.7.** This definition is a generalization of the cumulants in classical, free, Boolean and monotone probability theories.

We can prove the existence of cumulants.

**Theorem 4.8.** We assume the conditions (M1) and (M2) for a power associative convolution  $\square$ . Then cumulants of  $\square$  exist.

*Proof.* It is sufficient to prove that  $m_n(\mu^{\square N})$  is a polynomial of  $N$  by Proposition 4.4. Then the proof is the same one as given in [11], which we omit here.  $\square$

We discuss when the additivity of cumulants holds. In the proof of the following theorem, we assume the convolution is defined on  $\mathcal{P}_c$  to assure that moments determine a unique probability measure.

**Theorem 4.9.** Let  $\square$  be a power associative convolution defined on  $\mathcal{P}_c$  satisfying (M1) and (M2). Let  $r_n$  be the cumulants. Then the following conditions are equivalent.

- (1)  $r_n(\mu \square \nu) = r_n(\mu) + r_n(\nu)$  for all  $n$  and  $\mu, \nu \in \mathcal{P}_c$ .
- (2)  $\square$  is associative, commutative and  $P_n$  in (4.1) does not contain linear terms  $m_k(\mu)$  and  $m_k(\nu)$ ,  $1 \leq k \leq n-1$  for any  $n$ .

*Proof.* (1)  $\Rightarrow$  (2): The associativity and commutativity follow immediately since probability measures with compact supports are determined by the cumulants. By (M1) and (4.4) we obtain the identity

$$\begin{aligned} P_n(m_1(\mu), \dots, m_{n-1}(\mu), m_1(\nu), \dots, m_{n-1}(\nu)) \\ = Q_n(m_1(\mu), \dots, m_{n-1}(\mu)) + Q_n(m_1(\nu), \dots, m_{n-1}(\nu)) \\ + R_n(r_1(\mu) + r_1(\nu), \dots, r_{n-1}(\mu) + r_{n-1}(\nu)). \end{aligned}$$

By (C2'),  $Q_n$  and  $R_n$  do not contain linear terms.

(2)  $\Rightarrow$  (1): By (M1), (C2') and (4.4)

$$\begin{aligned} r_n(\mu \square \nu) &= m_n(\mu \square \nu) + Q_n(m_1(\mu \square \nu), \dots, m_{n-1}(\mu \square \nu)) \\ &= r_n(\mu) + r_n(\nu) + P_n(m_1(\mu), \dots, m_{n-1}(\mu), m_1(\nu), \dots, m_{n-1}(\nu)) \\ &\quad + R_n(r_1(\mu), \dots, r_{n-1}(\mu)) + R_n(r_1(\nu), \dots, r_{n-1}(\nu)) \\ &\quad + Q_n(m_1(\mu \square \nu), \dots, m_{n-1}(\mu \square \nu)). \end{aligned}$$

Therefore, there exists a polynomial  $U_n$  which does not contain linear terms such that  $r_n(\mu \square \nu) = r_n(\mu) + r_n(\nu) + U_n(r_1(\mu), \dots, r_{n-1}(\mu), r_1(\nu), \dots, r_{n-1}(\nu))$ . We replace  $\mu$  and  $\nu$  by  $\mu^{\square N}$  and  $\nu^{\square N}$ , respectively. We note that by the associativity and commutativity, it holds that  $r_n(\mu^{\square N} \square \nu^{\square N}) = r_n((\mu \square \nu)^{\square N}) = Nr_n(\mu \square \nu)$ . Then we obtain  $Nr_n(\mu \square \nu) = Nr_n(\mu) + Nr_n(\nu) + U_n(Nr_1(\mu), \dots, Nr_{n-1}(\mu), Nr_1(\nu), \dots, Nr_{n-1}(\nu))$ . This can be seen as an identity of polynomials of  $N$ ; therefore, we have  $U_n = 0$ .  $\square$

**Theorem 4.10.** *Let  $\square$  be a power associative convolution defined on  $\mathcal{P}_m$  satisfying (M1) and (M2). Let  $r_n$  be the cumulants.*

(1) (central limit theorem) *For  $\mu \in \mathcal{P}_m$  with  $m_1(\mu) = 0$  and  $m_2(\mu) = 1$ , we define  $\mu_N := (D_{\frac{1}{\sqrt{N}}} \mu)^{\square N}$ .*

*Then  $r_1(\mu_N) \rightarrow 0$ ,  $r_2(\mu_N) \rightarrow 1$  and  $r_n(\mu_N) \rightarrow 0$  as  $N \rightarrow \infty$  for any  $n \geq 3$ .*

(2) (Poisson's law of small numbers) *Let  $\{\mu^{(N)}\}$  be a sequence such that for any  $n \geq 1$   $Nm_n(\mu^{(N)}) \rightarrow \lambda > 0$  as  $N \rightarrow \infty$ . We define  $\mu_N := (\mu^{(N)})^{\square N}$ . Then  $r_n(\mu_N) \rightarrow \lambda$  as  $N \rightarrow \infty$  for any  $n \geq 1$ .*

## 5 Conditionally monotone cumulants

### 5.1 Additivity of conditionally monotone cumulants

We discuss the cumulants of conditionally monotone convolution based on Definition 4.6. Although conditionally monotone convolution is defined as a part of conditionally free convolution, conditionally free cumulants seem to be not suitable to look for the cumulants. For the second component of a pair of probability measures, we use monotone cumulants defined in [11]. We define conditionally monotone cumulants for the first component.

**Lemma 5.1.** *Let  $\mu \in \mathcal{P}_m$ . We define  $b_n = b_n(\mu)$  by  $H_\mu(z) = z \left( 1 + \sum_{n=1}^{\infty} \frac{b_n}{z^n} \right)$  in the sense of asymptotic expansion. Then  $b_n$  is of the form*

$$b_n = -m_n + W_n(m_1, \dots, m_{n-1}), \quad (5.1)$$

where  $W_n$  is a polynomial. Moreover,  $b_n(D_\lambda \mu) = \lambda^n b_n(\mu)$  for  $\lambda > 0$ . In particular,  $W_n$  does not contain a constant term and linear terms.

*Proof.* We have

$$\begin{aligned} H_\mu(z) &= z \frac{1}{1 + \sum_{k=1}^{\infty} \frac{m_k}{z^k}} \\ &= z \left( 1 - \sum_{k=1}^{\infty} \frac{m_k}{z^k} + \left( \sum_{k=1}^{\infty} \frac{m_k}{z^k} \right)^2 - \dots \right). \end{aligned} \quad (5.2)$$

Then we obtain the polynomial  $W_n$ . The last statement is obtained by the relation  $H_{D_\lambda \mu}(z) = \lambda H_\mu(\lambda^{-1} z)$ .  $\square$

**Lemma 5.2.** *For any  $n \geq 3$ , there exists a polynomial  $Y_n$  of  $2n - 3$  variables such that  $b_n(\mu_1 \triangleright_{\nu_2} \mu_2) = b_n(\mu_1) + b_n(\mu_2) + Y_n(b_1(\mu_1), \dots, b_{n-1}(\mu_1), b_1(\nu_2), \dots, b_{n-2}(\nu_2))$ . In addition,  $Y_n$  does not contain a constant term and linear terms. For  $n = 1$  and  $2$ , we have  $b_1(\mu_1 \triangleright_{\nu_2} \mu_2) = b_1(\mu_1) + b_1(\mu_2)$  and  $b_2(\mu_1 \triangleright_{\nu_2} \mu_2) = b_2(\mu_1) + b_2(\mu_2)$ .*

*Proof.* The additivity of  $b_1$  and  $b_2$  has been proved in Proposition 7.3. We have the equality

$$\begin{aligned}
H_{\mu_1 \triangleright_{\nu_2} \mu_2}(z) &= H_{\mu_1}(H_{\nu_2}(z)) + H_{\mu_2}(z) - H_{\nu_2}(z) \\
&= b_1(\mu_1) + \sum_{n=1}^{\infty} b_{n+1}(\mu_1) G_{\nu_2}(z)^n + z + b_1(\mu_2) + \sum_{n=1}^{\infty} \frac{b_{n+1}(\mu_2)}{z^n} \\
&= b_1(\mu_1) + \sum_{n=1}^{\infty} \frac{b_{n+1}(\mu_1)}{z^n} \left(1 + \sum_{k=1}^{\infty} \frac{m_k(\nu_2)}{z^k}\right)^n + z + b_1(\mu_2) + \sum_{n=1}^{\infty} \frac{b_{n+1}(\mu_2)}{z^n}.
\end{aligned} \tag{5.3}$$

By Lemma 5.1,  $m_k(\nu_2)$  has such a form as  $m_k(\nu_2) = -b_k(\nu_2) + X_k(b_1(\nu_2), \dots, b_{k-1}(\nu_2))$ , where  $X_k$  is a polynomial without a constant term and linear terms. Therefore, we have the conclusion.  $\square$

We now consider cumulants. For cumulants of the second component, we use monotone cumulants  $r_n(\nu)$  and its generating function  $A_\nu(z) = -\sum_{n=1}^{\infty} \frac{r_n(\nu)}{z^{n-1}}$ . In order to define conditionally monotone cumulants, we first give a formal discussion based on the formula (4.7). Let  $\{(\mu_t, \nu_t)\}_{t \geq 0}$  be a conditionally monotone convolution semigroup with  $(\mu_0, \nu_0) = (\delta_0, \delta_0)$ . Then  $\{\nu_t\}_{t \geq 0}$  is a monotone convolution semigroup with  $\nu_0 = \delta_0$ . We define  $\mu := \mu_1$  and  $\nu := \nu_1$ . By (7.7) we have  $H_{\mu_{s+t}} = H_{\mu_s} \circ H_{\nu_t} + H_{\mu_t} - H_{\nu_t}$ . We assume that  $m_n(\mu_t)$  is a differentiable function of  $t$ . Differentiating the equality by  $s$  formally and putting  $s = 0$ , we obtain

$$\frac{d}{dt} H_{\mu_t}(z) = A_{(\mu, \nu)}(H_{\nu_t}(z)), \tag{5.4}$$

where  $A_{(\mu, \nu)}(z)$  is defined by  $A_{(\mu, \nu)}(z) = \frac{d}{dt} H_{\mu_t}(z)|_{t=0}$ . We expand  $A_{(\mu, \nu)}(z)$  as a formal power series:

$$A_{(\mu, \nu)}(z) = -\sum_{n=1}^{\infty} \frac{r_n(\mu, \nu)}{z^{n-1}}. \tag{5.5}$$

(5.4) is equivalent to

$$\frac{d}{dt} G_{\mu_t}(z) = \sum_{n=1}^{\infty} r_n(\mu, \nu) G_{\mu_t}(z)^2 G_{\nu_t}(z)^{n-1} \tag{5.6}$$

as a formal power series. There exists a polynomial  $M_n$  for any  $n \geq 3$  such that

$$\begin{aligned}
\frac{d}{dt} m_n(\mu_t) &= r_n(\mu, \nu) + M_n(r_1(\mu, \nu), \dots, r_{n-1}(\mu, \nu), m_1(\mu_t), \dots, m_{n-1}(\mu_t), \\
&\quad m_1(\nu_t), \dots, m_{n-2}(\nu_t)).
\end{aligned} \tag{5.7}$$

It is easy to prove that  $M_1 = 0$  and  $M_2$  only depends on  $r_1(\mu, \nu)$  and  $m_1(\mu_t)$ . Since  $m_k(\nu_t)$  can be expressed by the polynomial of  $r_1(\nu), \dots, r_k(\nu)$ , we can prove inductively that  $m_n(\mu_t)$  is of the form

$$m_n(\mu_t) = r_n(\mu, \nu)t + t^2 C_n(t, r_1(\mu, \nu), \dots, r_{n-1}(\mu, \nu), r_1(\nu), \dots, r_{n-2}(\nu)), \tag{5.8}$$

where  $C_n$  is a polynomial for any  $n \geq 3$  and  $r_k(\mu)$  is the monotone  $k$ -th cumulant of  $\mu$ . It is easy to show that  $r_1(\mu, \nu) = m_1(\mu)$  and  $r_2(\mu, \nu) = \sigma^2(\mu) = m_2(\mu) - m_1(\mu)^2$ . Therefore,  $r_n(\mu, \nu)$  are expressed by polynomials of  $m_k(\mu)$  and  $m_k(\nu)$  ( $1 \leq k \leq n$ ). These relations can be generalized to any probability measures  $\mu, \nu \in \mathcal{P}_m$ .

**Definition 5.3.** For probability measures  $\mu, \nu \in \mathcal{P}_m$ , we define the conditionally monotone cumulants  $r_n(\mu, \nu)$  ( $n \geq 1$ ) by the equations  $m_n(\mu) = r_n(\mu, \nu) + C_n(1, r_1(\mu, \nu), \dots, r_{n-1}(\mu, \nu), r_1(\nu), \dots, r_{n-2}(\nu))$  for  $n \geq 3$ . For  $n = 1, 2$ , we define  $r_1(\mu, \nu) := m_1(\mu)$  and  $r_2(\mu, \nu) := m_2(\mu) - m_1(\mu)^2$ .

Now we prove the additivity of the cumulants. The proof is based on the relation of generating functions in Proposition 3.3. We note, however, that we can give a proof without the use of generating functions as shown in [11].

**Theorem 5.4.**  $r_n((\mu, \nu)^{\triangleright N}) = N r_n(\mu, \nu)$  holds for  $\mu, \nu \in \mathcal{P}_m$  and  $N \in \mathbb{N}$ .

*Proof.* We define  $\mu_N$  by  $(\mu_N, \nu^{\triangleright N}) = (\mu, \nu)^{\triangleright N}$ . We define

$$m_n(\nu, t) := \sum_{k=1}^n \sum_{1=i_0 < i_1 < \dots < i_{k-1} < i_k = n+1} \frac{t^k}{k!} \prod_{l=1}^k i_{l-1} r_{i_l - i_{l-1}}(\nu) \quad (5.9)$$

and

$$m_n(\mu, \nu, t) := r_n(\mu, \nu)t + t^2 C_n(t, r_1(\mu, \nu), \dots, r_{n-1}(\mu, \nu), r_1(\nu), \dots, r_{n-2}(\nu)), \quad (5.10)$$

which may not be moments of a probability measure for general  $t$ . It is noted that  $m_n(\mu, \nu, 1) = m_n(\mu)$  and  $m_n(\mu, \mu, t) = m_n(\mu, t)$ . (The latter equality comes from the relation  $(\mu, \mu) \triangleright (\nu, \nu) = (\mu \triangleright \nu, \mu \triangleright \nu)$ .) We shall show that  $m_n(\mu, \nu, N) = m_n(\mu_N, \nu^{\triangleright N}, 1) (= m_n(\mu_N))$  for any  $n, N \geq 1$ . We define formal power series  $A_{(\mu, \nu)}(z)$  by (5.5) and  $H_{(\mu, \nu)}(t, z) := \left( \sum_{n=0}^{\infty} \frac{m_n(\mu, \nu, t)}{z^{n+1}} \right)^{-1}$  and  $H_{\nu}(t, z) := \left( \sum_{n=0}^{\infty} \frac{m_n(\nu, t)}{z^{n+1}} \right)^{-1}$ . We note that the equality  $H_{\nu}(t+s, z) = H_{\nu}(t, H_{\nu}(s, z))$  holds as formal power series (see [11] for the proof). Now we show that

$$H_{(\mu, \nu)}(t+s, z) = H_{(\mu, \nu)}(t, H_{\nu}(s, z)) - H_{\nu}(s, z) + H_{(\mu, \nu)}(s, z). \quad (5.11)$$

We define  $K_s(t, z) := H_{(\mu, \nu)}(t, H_{\nu}(s, z)) - H_{\nu}(s, z) + H_{(\mu, \nu)}(s, z)$ . By the arguments given before Definition 5.3, we have

$$\frac{d}{dt} H_{(\mu, \nu)}(t, z) = A_{(\mu, \nu)}(H_{\nu}(t, z)). \quad (5.12)$$

It is clear that  $H_{(\mu, \nu)}(t+s, z)$  satisfies the differential equation  $\frac{d}{dt} H_{(\mu, \nu)}(t+s, z) = A_{(\mu, \nu)}(H_{\nu}(t+s, z))$  for fixed  $s$  with the initial value  $H_{(\mu, \nu)}(s, z)$ . On the other hand, by (5.12) we have  $\frac{d}{dt} K_s(t, z) = A_{(\mu, \nu)}(H_{\nu}(t, H_{\nu}(s, z))) = A_{(\mu, \nu)}(H_{\nu}(t+s, z))$ . Moreover, we have  $K_s(0, z) = H_{(\mu, \nu)}(s, z)$ . Therefore,  $H_{(\mu, \nu)}(t+s, z) = K_s(t, z)$ . Since the coefficients of  $z^k$  in the equality (5.11) are polynomials of  $t$  and  $s$ , we can substitute real numbers. Setting  $t = s = 1$ , we obtain  $m_n(\mu, \nu, 2) = m_n(\mu_2, \nu \triangleright \nu, 1) = m_n(\mu_2)$ . Inductively, we can show that  $m_n(\mu, \nu, N) = m_n(\mu_N)$ . By the (power) associativity of conditionally monotone convolution, we also obtain  $m_n(\mu, \nu, MN) = m_n(\mu_N, \nu^{\triangleright N}, M)$  for  $M, N \in \mathbb{N}$ . Then we have

$$\begin{aligned} & MN r_n(\mu, \nu) + M^2 N^2 C_n(MN, r_1(\mu, \nu), \dots, r_{n-1}(\mu, \nu), r_1(\nu), \dots, r_{n-2}(\nu)) \\ &= M r_n(\mu_N, \nu^{\triangleright N}) + M^2 C_n(M, r_1(\mu_N, \nu^{\triangleright N}), \dots, r_{n-1}(\mu_N, \nu^{\triangleright N}), r_1(\nu^{\triangleright N}), \dots, r_{n-2}(\nu^{\triangleright N})). \end{aligned} \quad (5.13)$$

Since the coefficients of  $M$  coincide, it holds that  $r_n(\mu_N, \nu^{\triangleright N}) = N r_n(\mu, \nu)$ .  $\square$

Before closing this section, we summarize the basic relations of generating functions:

$$\frac{d}{dt} H_{(\mu, \nu)}(t, z) = A_{(\mu, \nu)}(H_{\nu}(t, z)), \quad H_{(\mu, \nu)}(1, z) = H_{\mu}(z), \quad H_{(\mu, \nu)}(0, z) = z, \quad (5.14)$$

$$\frac{d}{dt} H_{\nu}(t, z) = A_{\nu}(H_{\nu}(t, z)), \quad H_{\nu}(1, z) = H_{\nu}(z), \quad H_{\nu}(0, z) = z. \quad (5.15)$$

These relations are parallel to

$$H_{\mu}(z) = z - \phi_{(\mu, \nu)}(H_{\nu}(z)), \quad (5.16)$$

$$H_{\nu}(z) = z - \phi_{\nu}(H_{\nu}(z)), \quad (5.17)$$

which are basic relations in conditionally free convolution. We note that monotone cumulants and Boolean cumulants are obtained as special cases of conditionally monotone cumulants:  $A_{(\mu, \mu)}$  is a generating function of monotone cumulants and  $A_{(\mu, \delta_0)}$  is a generating function of Boolean cumulants.

## 5.2 Moment-cumulant formula and monotone partitions

Next we show a combinatorial moment-cumulant formula for conditionally monotone convolution. Let  $\mathcal{LNC}(n)$  be the set of all linearly ordered non-crossing partitions defined by

$$\mathcal{LNC}(n) := \{(\pi, \lambda); \pi \in \mathcal{NC}(n), \lambda \text{ is a linear ordering of the blocks of } \pi\}. \quad (5.18)$$

We shall prove that the moment-cumulant formula for conditionally monotone convolution is given by

$$m_n(\mu) = \sum_{(\pi, \lambda) \in \mathcal{M}(n)} \frac{1}{|\pi|!} \left( \prod_{\substack{V \in \pi, \\ V: \text{ outer}}} r_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \pi, \\ V: \text{ inner}}} r_{|V|}(\nu) \right). \quad (5.19)$$

This formula is very parallel to conditionally free moment-cumulant formula which can be described as

$$m_n(\mu) = \sum_{(\pi, \lambda) \in \mathcal{LNC}(n)} \frac{1}{|\pi|!} \left( \prod_{\substack{V \in \pi, \\ V: \text{ outer}}} R_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \pi, \\ V: \text{ inner}}} R_{|V|}(\nu) \right) \quad (5.20)$$

if we impose the linear order structure. Of course the role of the linear order structure is trivial in (5.20), but it is crucial in (5.19). When we have defined conditionally monotone cumulants, we used the Taylor expansion of the equality  $\frac{d}{dt} H_{\mu_t}(z) = A_{(\mu, \nu)}(H_{\nu_t}(z))$ , which was useful to prove the additivity  $r_n((\mu, \nu)^{\triangleright N}) = N r_n(\mu, \nu)$ . When we try to obtain a combinatorial moment-cumulant formula, this equality is not useful and we use the following alternative one.

**Proposition 5.5.** *Let  $m_n(\mu, \nu, t)$ ,  $m_n(\mu, t)$ ,  $r_n(\mu)$  and  $r_n(\mu, \nu)$  be the quantities defined above. We have the recursive differential equations*

$$\begin{aligned} \frac{d}{dt} m_n(\mu, \nu, t) &= \sum_{k=0}^{n-1} (k+1) r_{n-k}(\nu) m_k(\mu, \nu, t) - \sum_{k=0}^{n-1} \sum_{l=0}^k r_{n-k}(\nu) m_l(\mu, \nu, t) m_{k-l}(\mu, \nu, t) \\ &\quad + \sum_{k=0}^{n-1} \sum_{l=0}^k r_{n-k}(\mu, \nu) m_l(\mu, \nu, t) m_{k-l}(\mu, \nu, t). \end{aligned} \quad (5.21)$$

**Remark 5.6.** When we put  $\mu = \nu$ , the above equation becomes  $\frac{d}{dt} m_n(\mu, t) = \sum_{k=0}^{n-1} (k+1) r_{n-k}(\mu) m_k(\mu, t)$  which has been obtained in the case of monotone convolution [11]. Moreover, we have  $\frac{d}{dt} m_n(\mu, \delta_0, t) = \sum_{k=0}^{n-1} \sum_{l=0}^k r_{n-k}(\mu, \delta_0) m_l(\mu, \delta_0, t) m_{k-l}(\mu, \delta_0, t)$  since  $r_n(\delta_0) = 0$ . In this case  $r_n(\mu, \delta_0)$  is an  $n$ -th Boolean cumulant. It might be interesting that this differential equation describes the structure of interval partitions.

*Proof.* We use the same notation as used in the subsection 5.1. Differentiating the equality (5.11) w.r.t.  $s$  and putting  $s = 0$  we obtain

$$\frac{\partial H_{(\mu, \nu)}}{\partial t}(t, z) = A_\nu(z) \frac{\partial H_{(\mu, \nu)}}{\partial z}(t, z) - A_\nu(z) + A_{(\mu, \nu)}(z). \quad (5.22)$$

We define  $G_{(\mu, \nu)}(t, z) := \frac{1}{H_{(\mu, \nu)}(t, z)}$ . Then we get the equality

$$\frac{\partial G_{(\mu, \nu)}}{\partial t}(t, z) = A_\nu(z) \frac{\partial G_{(\mu, \nu)}}{\partial z}(t, z) + G_{(\mu, \nu)}(t, z)^2 A_\nu(z) - G_{(\mu, \nu)}(t, z)^2 A_{(\mu, \nu)}(z). \quad (5.23)$$

(5.21) is obtained by the formal power series expansion of (5.23).  $\square$


We say a block  $V$  in a partition  $\pi \in \mathcal{P}(n)$  is of *interval type* if there exist  $j$  and  $k$  ( $1 \leq j \leq n$ ,  $0 \leq k \leq n-j$ ) such that  $V = \{j, j+1, \dots, j+k\}$ .

**Theorem 5.7.** *The moment-cumulant formula for conditionally monotone convolution is given by*

$$m_n(\mu) = \sum_{(\pi, \lambda) \in \mathcal{M}(n)} \frac{1}{|\pi|!} \left( \prod_{\substack{V \in \pi, \\ V: \text{ outer}}} r_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \pi, \\ V: \text{ inner}}} r_{|V|}(\nu) \right). \quad (5.24)$$

*Proof.* This proof gives the combinatorial meaning of the differential equations (5.21). We give a proof by induction. We assume that the formula (5.24), where  $m_n(\mu)$  is replaced by  $m_n(\mu, \nu, t)$  and  $\frac{1}{|\pi|!}$  is replaced by  $\frac{t^{|\pi|}}{|\pi|!}$ , is the case for  $1 \leq n \leq N$ . For a given monotone partition  $(\pi, \lambda) \in \mathcal{M}(N+1)$ , we

denote it by  $\pi = \{V_1 < \dots < V_{|\pi|}\}$ : this notation means that the indices of the blocks denote the linear ordering. We always use this notation in this proof. Note that  $V_{|\pi|}$  is a block of interval type. We assume  $|V_{|\pi|}| = N + 1 - k$  ( $0 \leq k \leq N$ ). We divide the summation into several cases focusing on the highest ordered block  $V_{|\pi|}$ . In this point of view, there are two cases where  $V_{|\pi|}$  is an outer block and  $V_{|\pi|}$  is an inner block. If  $V_{|\pi|}$  is an outer block,  $\pi$  needs to be of such a form as



where  $\sigma$  and  $\tau$  are arbitrary non-crossing partitions with  $\sigma \in \mathcal{NC}(l)$  and  $\tau \in \mathcal{NC}(k-l)$  ( $0 \leq l \leq k$ ). (When  $l = 0$ , we understand that  $\sigma = \emptyset$ .) Next we need to consider the linear orderings of  $\sigma$  and  $\tau$ . We note that a linear ordering of the blocks in  $\sigma \cup \tau$  is equivalent to distributing natural numbers  $\{1, \dots, |\sigma| + |\tau|\}$  to them such that  $\sigma$  and  $\tau$ , equipped with the natural numbers, become monotone partitions. If  $\sigma$  and  $\tau$  are once fixed, there are  $\frac{(|\sigma| + |\tau|)!}{|\sigma|!|\tau|!}$  possible ways to divide the set  $\{1, \dots, |\sigma| + |\tau|\}$  into two subsets  $C$  and  $D$  with  $|C| = |\sigma|$  and  $|D| = |\tau|$ . After such a division, we can choose linear orderings of  $\sigma$  from  $C$  and  $\tau$  from  $D$  independently with each other, and hence, we can take arbitrary  $\sigma \in \mathcal{M}(l)$  and  $\tau \in \mathcal{M}(k-l)$ . The above arguments imply that

$$\begin{aligned}
& \sum_{\substack{(\pi, \lambda) \in \mathcal{M}(N+1); \\ V_{|\pi|}: \text{outer}}} \frac{t^{|\pi|}}{|\pi|!} \left( \prod_{\substack{V \in \pi, \\ V: \text{outer}}} r_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \pi, \\ V: \text{inner}}} r_{|V|}(\nu) \right) \\
&= \sum_{k=0}^N \sum_{l=0}^k \sum_{\substack{(\sigma, \rho) \in \mathcal{M}(l) \\ (\tau, \lambda) \in \mathcal{M}(k-l)}} \frac{(|\sigma| + |\tau|)!}{|\sigma|!|\tau|!} \frac{t^{|\sigma| + |\tau| + 1} r_{n-k}(\mu, \nu)}{(|\sigma| + |\tau| + 1)!} \left( \prod_{\substack{V \in \sigma \cup \tau, \\ V: \text{outer}}} r_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \sigma \cup \tau, \\ V: \text{inner}}} r_{|V|}(\nu) \right) \quad (5.25) \\
&= \sum_{k=0}^N \sum_{l=0}^k r_{n-k}(\mu, \nu) \int_0^t m_l(\mu, \nu, s) m_{k-l}(\mu, \nu, s) ds,
\end{aligned}$$

where we have used the assumption of induction in the last line. Next we consider the case where  $V_{|\pi|}$  is an inner block. In this case, first we count all monotone partitions and later we remove the sums where  $V_{|\pi|}$  is an outer block. When  $|V_{|\pi|}| = N + 1 - k$ , there are  $k + 1$  ways to choose the position of  $V_{|\pi|}$ . Then we can take arbitrary monotone partition in  $\mathcal{M}(k)$  as if the highest block  $V_{|\pi|}$  were absent. (This idea is the same as used in [11].) The above arguments amount to

$$\begin{aligned}
& \sum_{\substack{(\pi, \lambda) \in \mathcal{M}(N+1); \\ V_{|\pi|}: \text{inner}}} \frac{t^{|\pi|}}{|\pi|!} \left( \prod_{\substack{V \in \pi, \\ V: \text{outer}}} r_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \pi, \\ V: \text{inner}}} r_{|V|}(\nu) \right) \\
&= \sum_{k=0}^N (k+1) r_{n-k}(\nu) \sum_{(\sigma, \rho) \in \mathcal{M}(k)} \frac{t^{|\sigma| + 1}}{(|\sigma| + 1)!} \left( \prod_{\substack{V \in \sigma, \\ V: \text{outer}}} r_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \sigma, \\ V: \text{inner}}} r_{|V|}(\nu) \right) \\
&\quad - \sum_{k=0}^N \sum_{l=0}^k \sum_{\substack{(\sigma, \rho) \in \mathcal{M}(l) \\ (\tau, \lambda) \in \mathcal{M}(k-l)}} \frac{(|\sigma| + |\tau|)!}{|\sigma|!|\tau|!} \frac{t^{|\sigma| + |\tau| + 1} r_{n-k}(\nu)}{(|\sigma| + |\tau| + 1)!} \left( \prod_{\substack{V \in \sigma \cup \tau, \\ V: \text{outer}}} r_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \sigma \cup \tau, \\ V: \text{inner}}} r_{|V|}(\nu) \right) \quad (5.26) \\
&= \sum_{k=0}^N (k+1) r_{n-k}(\nu) \int_0^t m_k(\nu, s) ds - \sum_{k=0}^N \sum_{l=0}^k r_{n-k}(\nu) \int_0^t m_l(\mu, \nu, s) m_{k-l}(\mu, \nu, s) ds.
\end{aligned}$$

By Proposition 5.5, we obtain

$$m_{N+1}(\mu, \nu, t) = \sum_{(\pi, \lambda) \in \mathcal{M}(N+1)} \frac{t^{|\pi|}}{|\pi|!} \left( \prod_{\substack{V \in \pi, \\ V: \text{outer}}} r_{|V|}(\mu, \nu) \right) \left( \prod_{\substack{V \in \pi, \\ V: \text{inner}}} r_{|V|}(\nu) \right). \quad (5.27)$$



## 6 Limit theorems of conditionally monotone convolution

We consider the central limit theorem and Poisson's law of small numbers of conditionally monotone convolution. Let  $\mu, \nu \in \mathcal{P}_m$  such that  $m_1(\mu) = m_1(\nu) = 0$  and  $m_2(\mu) = \alpha^2$  and  $m_2(\nu) = \beta^2$  with  $\alpha, \beta > 0$ . We define  $(\mu_N, \nu_N) := (D_{\frac{1}{\sqrt{N}}}\mu, D_{\frac{1}{\sqrt{N}}}\nu)^{\triangleright N}$ . By a simple calculation, we can show that  $r_1(\mu_N, \nu_N) = 0$ ,  $r_2(\mu_N, \nu_N) = \alpha^2$  and  $r_n(\mu_N, \nu_N) \rightarrow 0$  as  $N \rightarrow \infty$ . The second component is the usual monotone convolution. Therefore, there exist limit distributions  $(\nu_{\alpha, \beta}, \nu_\beta)$  at least in the sense of moments. The limit distributions are characterized by

$$\frac{d}{dt}H_{(\nu_{\alpha, \beta}, \nu_\beta)}(t, z) = -\frac{\alpha^2}{H_{\nu_\beta}(t, z)}, \quad H_{(\nu_{\alpha, \beta}, \nu_\beta)}(1, z) = H_{\nu_{\alpha, \beta}}(z), \quad (6.1)$$

$$\frac{d}{dt}H_{\nu_\beta}(t, z) = -\frac{\beta^2}{H_{\nu_\beta}(t, z)}, \quad H_{\nu_\beta}(1, z) = H_{\nu_\beta}(z). \quad (6.2)$$

We obtain  $H_{\nu_\beta}(t, z) = \sqrt{z^2 - 2\beta^2 t}$  from (6.2). Setting  $t = 1$ , we obtain the limit measure  $\nu_\beta(dx) = \frac{1}{\pi\sqrt{2\beta^2 - x^2}}dx$ ,  $x \in (-\sqrt{2}\beta, \sqrt{2}\beta)$ . For the first component, we obtain  $H_{(\nu_{\alpha, \beta}, \nu_\beta)}(1, z) = (1 - \frac{\alpha^2}{\beta^2})z + \frac{\alpha^2}{\beta^2}\sqrt{z^2 - 2\beta^2}$ , and hence, we have

$$G_{\nu_{\alpha, \beta}}(z) = \frac{\alpha^2\sqrt{z^2 - 2\beta^2} + (\alpha^2 - \beta^2)z}{(2\alpha^2 - \beta^2)z^2 - 2\alpha^4}. \quad (6.3)$$

This is the Stieltjes transform of the Kesten distribution. If we use the  $t$ -transformation, the limit distribution can be written by  $\mathcal{U}_{\alpha^2/\beta^2}(\nu_\beta)$ . The limit distribution is compactly supported, and then the convergence holds in the sense of weak convergence. We note that  $\nu_{\beta, \beta} = \nu_\beta$ .

Next we consider Poisson's law of small numbers in the setting of triangular arrays. Let  $\mu^{(N)}, \nu^{(N)} \in \mathcal{P}_m$  such that  $Nm_n(\mu^{(N)}) \rightarrow \lambda > 0$  and  $Nm_n(\nu^{(N)}) \rightarrow \rho > 0$  as  $N \rightarrow \infty$  for any  $n \geq 1$ . We define  $(\mu_N, \nu_N) := (\mu^{(N)}, \nu^{(N)})^{\triangleright N}$ . It is not difficult to prove that  $r_n(\mu_N, \nu_N) \rightarrow \lambda$  as  $N \rightarrow \infty$  for any  $n \geq 1$ . It is known that  $\nu_N$  converges weakly to the monotone Poisson distribution  $p_\rho$  with parameter  $\rho$ , which is characterized by the differential equation

$$\frac{d}{dt}H_{p_\rho}(t, z) = \frac{\rho H_{p_\rho}(t, z)}{1 - H_{p_\rho}(t, z)}, \quad H_{p_\rho}(1, z) = H_{p_\rho}(z), \quad (6.4)$$

where  $H_{p_\rho}(1, z) = H_{p_\rho}(z)$ . We denote the limit distribution  $\lim_{N \rightarrow \infty} \mu_N$  by  $p_{\lambda, \rho}$  in the sense of moments. Then  $p_{\lambda, \rho}$  is characterized by the differential equation

$$\frac{d}{dt}H_{(p_{\lambda, \rho}, p_\rho)}(t, z) = \frac{\lambda H_{p_\rho}(t, z)}{1 - H_{p_\rho}(t, z)}, \quad H_{(p_{\lambda, \rho}, p_\rho)}(1, z) = H_{p_{\lambda, \rho}}(z). \quad (6.5)$$

By (6.4) and (6.5) we have  $\frac{d}{dt}H_{(p_{\lambda, \rho}, p_\rho)}(t, z) = \frac{\lambda}{\rho} \frac{d}{dt}H_{p_\rho}(t, z)$ , which implies

$$H_{p_{\lambda, \rho}}(z) = \left(1 - \frac{\lambda}{\rho}\right)z + \frac{\lambda}{\rho}H_{p_\rho}(z). \quad (6.6)$$

It is not difficult to see that  $p_{\lambda, \rho}$  has a compact support. Therefore,  $\mu_N$  converges weakly to  $p_{\lambda, \rho}$ . Also in this case,  $p_{\lambda, \rho}$  can be written as  $p_{\lambda, \rho} = \mathcal{U}_{\lambda/\rho}(p_\rho)$ .

## 7 Convolutions arising from conditionally monotone convolution

Conditionally monotone convolution generates monotone convolution and Boolean convolution in the left component:

$$(\mu, \mu) \triangleright (\nu, \nu) = (\mu \triangleright \nu, \mu \triangleright \nu), \quad (7.1)$$

$$(\mu, \delta_0) \triangleright (\nu, \delta_0) = (\mu \uplus \nu, \delta_0), \quad (7.2)$$

which are very parallel relations to (3.5) and (3.6). For a map  $T: \mathcal{P} \rightarrow \mathcal{P}$ , one can define a new convolution  $\boxplus_T$  by

$$(\mu \boxplus_T \nu, T\mu \boxplus T\nu) = (\mu, T\mu) \boxplus (\nu, T\nu). \quad (7.3)$$

A problem of Bożejko is to find all maps  $T: \mathcal{P} \rightarrow \mathcal{P}$  such that

$$T(\mu \boxplus_T \nu) = T\mu \boxplus T\nu. \quad (7.4)$$

If such  $T$  is found, then the new convolution is associative and commutative. Moreover, this is a generalization of free and Boolean convolutions. This problem has been considered in [7, 15, 16, 29], in which several such convolutions have been found. Motivated by these works, we consider the following type of convolution:

$$(\mu \triangleright_T \nu, T\nu) = (\mu, \delta_0) \boxplus (\nu, T\nu). \quad (7.5)$$

This relation is parallel to (7.3) in terms of conditionally monotone convolution:

$$(\mu \triangleright_T \nu, T\mu \triangleright T\nu) = (\mu, T\mu) \triangleright (\nu, T\nu). \quad (7.6)$$

Clearly, this convolution includes Boolean and monotone convolutions if we take  $T$  as  $T\mu = \delta_0$  for all  $\mu$  and  $T = \text{Id}$ , respectively.

The convolution may not be associative. Now we give a characterization when the convolution  $\triangleright_T$  becomes associative.

**Proposition 7.1.** (1)  $\triangleright_T$  is characterized by the equality

$$H_{\mu \triangleright_T \nu} = H_\mu \circ H_{T\nu} + H_\nu - H_{T\nu}. \quad (7.7)$$

(2) The convolution  $\triangleright_T$  becomes associative if and only if

$$T(\mu \triangleright_T \nu) = T\mu \triangleright T\nu \quad (7.8)$$

for all  $\mu$  and  $\nu$ .

**Remark 7.2.** (1) In many cases  $T$  is only defined in a subset of  $\mathcal{P}$  such as  $\mathcal{P}_m$ . In such a case, the above Proposition holds if the subset is closed under the convolution  $\triangleright_T$ . Hereafter, we often state results about the convolution  $\triangleright_T$  only for  $T: \mathcal{P} \rightarrow \mathcal{P}$  if such a generalization of the domain of  $T$  is trivial.

(2) It seems to be not known whether the condition (7.4) is necessary condition for the associativity of  $\boxplus_T$ .

*Proof.* (1) The proof is easy.

(2) First we calculate  $H_{(\mu \triangleright_T \nu) \triangleright_T \lambda}$ .

$$\begin{aligned} H_{(\mu \triangleright_T \nu) \triangleright_T \lambda} &= H_{\mu \triangleright_T \nu} \circ H_{T\lambda} + H_\lambda - H_{T\lambda} \\ &= H_\mu \circ H_{T\nu} \circ H_{T\lambda} + H_\nu \circ H_{T\lambda} - H_{T\nu} \circ H_{T\lambda} + H_\lambda - H_{T\lambda}. \end{aligned} \quad (7.9)$$

$H_{\mu \triangleright_T (\nu \triangleright_T \lambda)}$  is calculated as follows.

$$\begin{aligned} H_{\mu \triangleright_T (\nu \triangleright_T \lambda)} &= H_\mu \circ H_{T(\nu \triangleright_T \lambda)} + H_{\nu \triangleright_T \lambda} - H_{T(\nu \triangleright_T \lambda)} \\ &= H_\mu \circ H_{T(\nu \triangleright_T \lambda)} + H_\nu \circ H_{T\lambda} + H_\lambda - H_{T\lambda} - H_{T(\nu \triangleright_T \lambda)}. \end{aligned} \quad (7.10)$$

Then the associativity of the convolution implies that  $H_\mu \circ H_{T\nu} \circ H_{T\lambda} - H_\mu \circ H_{T(\nu \triangleright_T \lambda)}$  only depends on  $\nu$  and  $\lambda$ . If  $H_{T\nu} \circ H_{T\lambda}$  were not equal to  $H_{T(\nu \triangleright_T \lambda)}$  for some  $\nu$  and  $\lambda$ , then there would exist  $w \in \mathbb{C} \setminus \mathbb{R}$  such that  $H_{T\nu} \circ H_{T\lambda}(w) \neq H_{T(\nu \triangleright_T \lambda)}(w)$ . In this case,  $H_\mu \circ H_{T\nu} \circ H_{T\lambda}(w) - H_\mu \circ H_{T(\nu \triangleright_T \lambda)}(w)$  clearly depends on  $\mu$ , which is a contradiction. Therefore, we conclude that

$$H_{T\nu} \circ H_{T\lambda} = H_{T(\nu \triangleright_T \lambda)} \quad (7.11)$$

for all  $\nu, \lambda \in \mathcal{P}$ . Conversely, if (7.11) holds, it is not difficult to see that the convolution is associative.  $\square$

We show the additivity of mean and variance, which will be used in the proof of Theorem 8.2.

**Proposition 7.3.** *Let  $T: \mathcal{P} \rightarrow \mathcal{P}$  be an arbitrary map. Then we have the following properties.*

- (1)  $\mathcal{P}^2$  is closed under the convolution  $\triangleright_T$ .
- (2)  $m(\mu)$  and  $\sigma^2(\mu)$  are additive w.r.t. the convolution  $\triangleright_T$  considered in  $\mathcal{P}^2$ :

$$m(\mu \triangleright_T \nu) = m(\mu) + m(\nu), \quad (7.12)$$

$$\sigma^2(\mu \triangleright_T \nu) = \sigma^2(\mu) + \sigma^2(\nu). \quad (7.13)$$

*Proof.* We use the notation

$$H_\mu(z) = -m(\mu) + z + \int \frac{1}{x-z} d\rho_\mu(x). \quad (7.14)$$

Then we have

$$\begin{aligned} H_{\mu \triangleright_T \nu}(z) &= H_\mu \circ H_{T\nu}(z) + H_\nu(z) - H_{T\nu}(z) \\ &= -m(\mu) + H_{T\nu}(z) + \int_{\mathbb{R}} \frac{1}{x - H_{T\nu}(z)} d\rho_\mu(x) + H_\nu(z) - H_{T\nu}(z) \\ &= -m(\mu) + H_\nu(z) + \int_{\mathbb{R}} \frac{1}{x-z} d\left( \int_{\mathbb{R}} (T\nu)_y(x) d\rho_\mu(y) \right) \\ &= -m(\mu) - m(\nu) + z + \int_{\mathbb{R}} \frac{1}{x-z} d\left( \int_{\mathbb{R}} (T\nu)_y(x) d\rho_\mu(y) + \rho_\nu(x) \right), \end{aligned} \quad (7.15)$$

where  $\lambda_y \in \mathcal{P}$  is defined by  $H_{\lambda_y} = H_\lambda - y$  for  $\lambda \in \mathcal{P}$ . By Lemma 2.16,  $\mathcal{P}^2$  is closed under the convolution; moreover, the mean and the variance are additive.  $\square$

Thus,  $\mathcal{P}^2$  is closed under the convolution  $\triangleright_T$  for arbitrary map  $T$ . On the contrary, the closedness of  $\mathcal{P}_+$  and  $\mathcal{P}_{\text{sym}}$  under the convolution  $\triangleright_T$  does not hold without any assumptions. We show the necessary and sufficient conditions.

**Proposition 7.4.** (1) *The following two conditions are equivalent.*

- (1a)  $T(\mathcal{P}_+) \subset \mathcal{P}_+$ ,
- (1b)  $\mu \triangleright_T \nu \in \mathcal{P}_+$  for all  $\mu, \nu \in \mathcal{P}_+$ .

(2) *The following two conditions are equivalent.*

- (2a)  $T(\mathcal{P}_{\text{sym}}) \subset \mathcal{P}_{\text{sym}}$ ,
- (2b)  $\mu \triangleright_T \nu \in \mathcal{P}_{\text{sym}}$  for all  $\mu, \nu \in \mathcal{P}_{\text{sym}}$ .

*Proof.* (1) We assume (1a). By Lemma 2.3,  $H_\lambda$  is analytic in  $\mathbb{C} \setminus [0, \infty)$  and  $H_\lambda < 0$  in  $(-\infty, 0)$  for  $\lambda = \mu, \nu, T\nu$ . Then the composition  $H_\mu \circ H_{T\nu}$  is analytic in  $\mathbb{C} \setminus [0, \infty)$ , and hence,  $H_{\mu \triangleright_T \nu}$  is also analytic in the same region. This implies the condition (1) in Lemma 2.3. We use the notation

$$H_\mu(z) = b_\mu + z + \int_{\mathbb{R}} \frac{1+xz}{x-z} d\eta_\mu(x) \quad (7.16)$$

and the similar notation for  $H_\nu$ . We put  $g_\mu(z) := \int_{\mathbb{R}} \frac{1+xz}{x-z} d\eta_\mu(x)$ . By Proposition 7.1, we have

$$\begin{aligned} H_{\mu \triangleright_{\mathcal{T}} \nu}(z) &= H_\mu \circ H_{T\nu}(z) + H_\nu(z) - H_{T\nu}(z) \\ &= b_\mu + g_\mu(H_{T\nu}(z)) + H_\nu(z). \end{aligned} \quad (7.17)$$

Since  $g_\mu$  and  $H_{T\nu}$  are nondecreasing,  $g_\mu \circ H_{T\nu}$  is also nondecreasing. Then  $b_\mu + g_\mu \circ H_{T\nu}(-0) \leq b_\mu + g_\mu(-0) = H_\mu(-0) \leq 0$ . Therefore, we obtain  $H_{\mu \triangleright_{\mathcal{T}} \nu}(-0) \leq 0$ .

Next we assume (1b). We shall prove the fact by *reductio ad absurdum*; we assume that there exists  $\nu \in \mathcal{P}_+$  such that  $T\nu \notin \mathcal{P}_+$ . In the notation (7.15), we have

$$\begin{aligned} H_{\mu \triangleright_{\mathcal{T}} \nu}(z) &= H_\mu \circ H_{T\nu}(z) + H_\nu(z) - H_{T\nu}(z) \\ &= -m(\mu) + H_\nu(z) + \int_{\mathbb{R}} \frac{1}{x-z} d\left( \int_{\mathbb{R}} (T\nu)_y(x) d\rho_\mu(y) \right) \\ &= -m(\mu) + H_\nu(z) - G_{\rho_\mu \triangleright_{\mathcal{T}} T\nu}(z) \end{aligned} \quad (7.18)$$

for all  $\mu \in \mathcal{P}^2$ , where we defined  $\rho_\mu \triangleright_{\mathcal{T}} T\nu$  by the affinity in the left component of monotone convolution. We can construct  $\mu \in \mathcal{P}_+$  such that  $a(\rho_\mu) = 0$ . Then from (3) of Lemma 2.2 we have  $a(\rho_\mu \triangleright_{\mathcal{T}} T\nu) \leq a(T\nu) < 0$ , which means that  $G_{\rho_\mu \triangleright_{\mathcal{T}} T\nu}$  is not analytic in  $\mathbb{C} \setminus [0, \infty)$ . By assumption, both  $H_{\mu \triangleright_{\mathcal{T}} \nu}(z)$  and  $H_\nu(z)$  are analytic in  $\mathbb{C} \setminus [0, \infty)$ , which is a contradiction to the equality (7.18) (we note that Lemma 2.2 is applicable for any positive finite measure).

(2) We assume (2a). Since symmetric probability measures are characterized by  $H_\mu(-z) = -H_\mu(z)$  for  $z \in \mathbb{C} \setminus \mathbb{R}$ , the proof is not difficult.

Conversely, we assume (2b). We take  $\mu$  to be the arcsine law with mean 0 and variance 1. Clearly,  $\mu \in \mathcal{P}_{\text{sym}}$ . By assumption,  $H_{\mu \triangleright_{\mathcal{T}} \nu}(-z) = -H_{\mu \triangleright_{\mathcal{T}} \nu}(z)$  for all  $\nu \in \mathcal{P}_{\text{sym}}$ . Then (7.7) implies that

$$\sqrt{H_{T\nu}(z)^2 - 2} - \sqrt{H_{T\nu}(-z)^2 - 2} = H_{T\nu}(z) + H_{T\nu}(-z) \quad (7.19)$$

for  $z \in \mathbb{C}_+$ . After some calculations we obtain  $H_{T\nu}(-z) = -H_{T\nu}(z)$ , which means  $T\nu \in \mathcal{P}_{\text{sym}}$ .  $\square$

**Remark 7.5.** The above property is a generalization of the Boolean and monotone cases. In the cases of Boolean and monotone, moreover, we can show that  $\nu^{\square n} \in \mathcal{P}_+$  implies  $\nu \in \mathcal{P}_+$  ( $\square$  is Boolean or monotone convolution).

Sometimes the limit distribution of Poisson's law of small numbers concerning a deformed convolution does not belong to  $\mathcal{P}_+$  [15]. We can give a sufficient condition under which the limit distribution belongs to  $\mathcal{P}_+$  in the case of the convolution  $\triangleright_{\mathcal{T}}$ .

**Corollary 7.6.** (1) We assume that  $T(\mathcal{P}_+) \subset \mathcal{P}_+$ . If Poisson's law of small numbers holds, the limit distribution belongs to  $\mathcal{P}_+$ .

(2) We assume that  $T(\mathcal{P}_{\text{sym}}) \subset \mathcal{P}_{\text{sym}}$ . If the central limit theorem holds, the limit distribution belongs to  $\mathcal{P}_{\text{sym}}$ .

**Remark 7.7.** Poisson's law of small numbers and the central limit theorem mean the statements as given in Theorem 11.6.

*Proof.* If we take  $\mu^{(N)} := (1 - \frac{\lambda}{N})\delta_0 + \frac{\lambda}{N}\delta_1 \in \mathcal{P}_+$ , the limit distribution  $\lim_{N \rightarrow \infty} (\mu^{(N)})^{\triangleright_{\mathcal{T}} N} \in \mathcal{P}_+$  by Lemma 2.6. For the central limit theorem, we take  $\mu := \frac{1}{2}(\delta_{-1} + \delta_1)$ . Then  $(D_{\frac{1}{\sqrt{N}}}\mu)^{\triangleright_{\mathcal{T}} N} \in \mathcal{P}_{\text{sym}}$  and the limit distribution also belongs to  $\mathcal{P}_{\text{sym}}$  by Lemma 2.6.  $\square$

## 8 $V_{t,u,a}$ -transformation

We introduce  $V_{t,u,a}$ -transformation and prove that the transformation satisfies the condition (7.8). Moreover, this family of transformations includes the generalized  $t$ -transformation [16] and the  $V_a$ -transformation [15].

Let  $f: \mathcal{Q} \rightarrow \mathbb{R}$ , where  $\mathcal{Q}$  is a subset of  $\mathcal{P}$ . Typically  $\mathcal{Q}$  is chosen to be  $\mathcal{P}^2$ ,  $\mathcal{P}_m$  or  $\mathcal{P}_c$ . Motivated by the generalized  $t$ -transformation in [16] and  $V_a$ -transformation in [15], we look for a transform  $V_{t,f}$  of the form

$$\mu \mapsto V_{t,f}\mu, \quad H_{V_{t,f}\mu}(z) = tH_\mu(z) + (1-t)z + f(\mu). \quad (8.1)$$

If  $f(\mu) = (t-u)m(\mu)$ , this is the same as the generalized  $t$ -transformation. If  $t = 1$  and  $f(\mu) = ar_\mu(2)$ , this is the same as  $V_a$ -transformation ( $r_\mu(2)$  is the second free cumulant of  $\mu$ , which is also equal to the variance).

**Lemma 8.1.** *Assume that  $\mathcal{Q}$  is closed under the convolution  $\triangleright_{V_{t,f}}$ .  $V_{t,f}$  satisfies the associativity condition (7.8) considered in  $\mathcal{Q}$  if and only if  $f(\mu \triangleright_{V_{t,f}} \nu) = f(\mu) + f(\nu)$  for all  $\mu \in \mathcal{Q}$ .*

*Proof.* We denote  $\triangleright_{V_{t,f}}$  by  $\triangleright_{t,f}$  for simplicity. By (7.7) we obtain

$$\begin{aligned} H_{V_{t,f}(\mu \triangleright_{t,f} \nu)}(z) &= tH_{\mu \triangleright_{t,f} \nu}(z) + (1-t)z + f(\mu \triangleright_{t,f} \nu) \\ &= tH_\mu \circ H_{V_{t,f}\nu}(z) + tH_\nu(z) - tH_{V_{t,f}\nu}(z) + (1-t)z + f(\mu \triangleright_{t,f} \nu) \\ &= tH_\mu \circ H_{V_{t,f}\nu}(z) + tH_\nu(z) + (1-t)z + f(\nu) - tH_{V_{t,f}\nu}(z) + f(\mu \triangleright_{t,f} \nu) - f(\nu) \\ &= tH_\mu \circ H_{V_{t,f}\nu}(z) + (1-t)H_{V_{t,f}\nu}(z) + f(\mu \triangleright_{t,f} \nu) - f(\nu). \end{aligned} \quad (8.2)$$

On the other hand, we have

$$\begin{aligned} H_{V_{t,f}\mu \triangleright_{t,f} \nu}(z) &= H_{V_{t,f}\mu} \circ H_{V_{t,f}\nu}(z) \\ &= tH_\mu \circ H_{V_{t,f}\nu}(z) + (1-t)H_{V_{t,f}\nu}(z) + f(\mu). \end{aligned} \quad (8.3)$$

Therefore, the associativity condition (7.8) is equivalent to  $f(\mu \triangleright_{t,f} \nu) = f(\mu) + f(\nu)$ .  $\square$

We define  $V_{t,u,a}$ -transformation by letting

$$f(\mu) = f_{t,u,a}(\mu) = (t-u)m(\mu) + a\sigma^2(\mu). \quad (8.4)$$

More clearly, we define

$$H_{V_{t,u,a}\mu}(z) = tH_\mu(z) + (1-t)z + (t-u)m(\mu) + a\sigma^2(\mu). \quad (8.5)$$

We shall prove that  $V_{t,u,a}$ -transformation gives an associative convolution defined on  $\mathcal{P}^2$ . It may be interesting to consider  $f$  including higher order moments, which we do not treat now. We use the notation  $\triangleright_{t,u,a}$  for the convolution defined by  $V_{t,u,a}$ .

**Theorem 8.2.** *The convolution  $\triangleright_{t,u,a}$  defined on  $\mathcal{P}^2$  is associative.*

*Proof.* This fact follows from Lemma 8.1 and Proposition 7.3.  $\square$

In order to calculate the inverse transformation of  $V_{t,u,a}$ , we show the following facts.

**Lemma 8.3.** *We have the following equalities.*

$$m(V_{t,u,a}\mu) = um(\mu) - a\sigma^2(\mu), \quad (8.6)$$

$$\sigma^2(V_{t,u,a}\mu) = t\sigma^2(\mu). \quad (8.7)$$

*Proof.* We use the same notation as in Theorem 8.2. We have

$$H_{V_{t,u,a}\mu}(z) = -um(\mu) + a\sigma^2(\mu) + z + t \int \frac{1}{x-z} d\rho_\mu(x), \quad (8.8)$$

which implies the conclusion by Lemma 2.16.  $\square$

**Proposition 8.4.** *We have the following equality:*

$$V_{t',u',a'}V_{t,u,a} = V_{t't,u'u,a'+a't} \text{ for } t \geq 0, \ u, \ a \in \mathbb{R}. \quad (8.9)$$

*In particular, we have*

$$V_{t,u,a}^{-1} = V_{t^{-1},u^{-1},-\frac{a}{tu}} \text{ for } t > 0, \ u \neq 0, \ a \in \mathbb{R}. \quad (8.10)$$

**Proposition 8.5.** (1)  $\mathcal{P}_+ \cap \mathcal{P}^2$  is closed under the convolution  $\triangleright_{t,u,a}$  if and only if  $u \geq t$  and  $a = 0$ .  
(2)  $\mathcal{P}_{\text{sym}} \cap \mathcal{P}^2$  is closed under the convolution  $\triangleright_{t,u,a}$  if and only if  $a = 0$ .

*Proof.* These facts are easy consequences of Proposition 7.4.  $\square$

**Example 8.6.** (1) Oravecz has introduced Fermi convolution  $\bullet$  in [28]. Oravecz has mentioned the relation between Fermi convolution and conditionally free convolution:

$$(\mu \bullet \nu, \delta_{m(\mu)} \boxplus \delta_{m(\nu)}) = (\mu, \delta_{m(\mu)}) \boxplus (\nu, \delta_{m(\nu)}), \quad (8.11)$$

where  $m(\mu)$  denotes the mean of  $\mu$ . We can easily extend the Fermi convolution to the convolution coming from the map  $F_u$  defined by  $F_u \mu = \delta_{um(\mu)}$ . We have  $V_{0,u,0} = F_u$ ; therefore, we obtain an associative convolution  $\triangleright_{F_u}$  related to Fermi convolution defined by

$$(\mu \triangleright_{F_u} \nu, \delta_{um(\mu)+um(\nu)}) = (\mu, \delta_{um(\mu)}) \triangleright (\nu, \delta_{um(\nu)}). \quad (8.12)$$

(2)  $\mathcal{U}_t := V_{t,t,0}$  is called the  $t$ -transformation. We obtain an associative convolution  $\triangleright_t$  arising from  $\mathcal{U}_t$  defined by

$$(\mu \triangleright_t \nu, \mathcal{U}_t(\mu) \triangleright \mathcal{U}_t(\nu)) = (\mu, \mathcal{U}_t(\mu)) \triangleright (\nu, \mathcal{U}_t(\nu)). \quad (8.13)$$

We note that the  $t$ -transformation gives an interpolation between Boolean and monotone convolutions in the present situation: monotone convolution is obtained when  $t = 1$  and Boolean convolution is obtained when we take the limit  $t \rightarrow 0$ .

(3)  $V_a$ -transformation is equal to  $V_{1,1,a}$ .

In the following we discuss the meaning of these results. It is known that the  $t$ -transformation  $\mathcal{U}_t$  ( $t > 0$ ) satisfies the condition (7.4), which gives a family of new convolutions  $\boxplus_{\mathcal{U}_t}$  [7]. These convolutions can also be written as

$$\mu \boxplus_{\mathcal{U}_t} \nu = \mathcal{U}_{1/t}(\mathcal{U}_t(\mu) \boxplus \mathcal{U}_t(\nu)). \quad (8.14)$$

Apart from the context of conditionally free convolution, it seems interesting to study the deformations of Boolean and classical convolutions defined by the right hand side of (8.14), with  $\boxplus$  replaced by  $\uplus$  and  $*$ , respectively. The new convolutions have been studied in [8]. By definition, the deformed convolutions are associative and commutative.

We can also define the same deformations in the monotone case. The results in this section show that the deformation has a natural meaning in terms of conditionally monotone convolution as in the case of free convolution (cf. (7.3) and (7.6)). Of course the above discussion is meaningful for any  $T$  which is invertible such as some class of  $\mathbf{t}$ -transformations.

## 9 Deformations related to monotone infinitely divisible distributions

Krystek and Wojakowski have introduced a deformation connected to a  $\boxplus$ -infinitely divisible distribution in [15], which we explain now. For a  $\boxplus$ -infinitely divisible distribution  $\varphi$  with compact support, there corresponds a unique weakly continuous  $\boxplus$ -convolution semigroup  $\{\varphi_t\}_{t \geq 0}$  with  $\varphi_0 = \delta_0$  and  $\varphi_1 = \varphi$ . Define a transformation  $\Phi_t^\varphi$  by

$$\Phi_t^\varphi \mu = \varphi_{\sigma^2(\mu)t}. \quad (9.1)$$

This map satisfies the condition (7.4).

We introduce the monotone analog of  $\Phi_t^\varphi$ .

**Definition 9.1.** For a  $\triangleright$ -infinitely divisible distribution  $\xi \in \mathcal{P}^2$ , let  $\{\xi_t\}_{t \geq 0}$  be the corresponding weakly continuous  $\triangleright$ -convolution semigroup with  $\xi_0 = \delta_0$  and  $\xi_1 = \xi$ . Let  $f: \mathcal{P}^2 \rightarrow \mathbb{R}$ . We define a transformation  $\Xi_f^\xi$  by

$$\Xi_f^\xi \mu := \xi_{f(\mu)}. \quad (9.2)$$

**Lemma 9.2.**  $\Xi_f^\xi$  satisfies the associativity condition (7.8) in  $\mathcal{P}^2$  if and only if  $f(\mu \triangleright_{\Xi_f^\xi} \nu) = f(\mu) + f(\nu)$  for all  $\mu, \nu \in \mathcal{P}^2$ .

*Proof.* This fact follows from the equality  $\Xi_f^\xi \mu \triangleright \Xi_f^\xi \nu = \xi_{f(\mu)+f(\nu)}$ .  $\square$

**Theorem 9.3.** The map  $\Xi_t^\xi$  ( $t \geq 0$ ) defined by  $f(\mu) = t\sigma^2(\mu)$  satisfies the condition (7.8).

*Proof.* The fact follows from Proposition 7.3.  $\square$

**Remark 9.4.** (1) If  $\xi = \delta_a$ , the map  $f_{s,t}(\mu) = -sm(\mu) + t\sigma^2(\mu)$  ( $s, t \in \mathbb{R}$ ) is also possible. In this case we have  $\Xi_{f_{u,a}}^{\delta_1} = V_{0,u,a}$ .

(2) It may be interesting to consider higher moments than two, which we do not consider now.

**Proposition 9.5.** (1)  $\mathcal{P}_+ \cap \mathcal{P}^2$  is closed under the convolution  $\triangleright_{\Xi_t^\xi}$  if and only if  $\xi \in \mathcal{P}_+ \cap \mathcal{P}^2$ .  
(2) We assume that  $\xi \in \mathcal{P}_c$ . Then  $\mathcal{P}_{\text{sym}} \cap \mathcal{P}_c$  is closed under the convolution  $\triangleright_{\Xi_t^\xi}$  if and only if  $\xi \in \mathcal{P}_{\text{sym}} \cap \mathcal{P}_c$ .

*Proof.* This is an immediate consequence of Lemma 2.4, Lemma 2.5 and Proposition 7.4.  $\square$

## 10 Cumulants of deformed convolutions

We define  $r_n^T(\mu) := r_n(\mu, T\mu)$ . We show that  $r_n^T(\mu)$  is suitable for the cumulants of the convolution  $\triangleright_T$ .

**Proposition 10.1.** We assume that there exists a polynomial  $V_n$  of  $n+1$  variables, which does not contain a constant term, such that

$$m_n(T\mu) = V_n(m_1(\mu), \dots, m_{n+1}(\mu)) \quad (10.1)$$

for any  $n \geq 1$ . Then the conditions (M1) and (M2) hold for the convolution  $\triangleright_T$ .

*Proof.* By (5.1) and (10.1), we obtain (M1). (M2) follows from the facts that both  $W_n$  and  $Y_n$  do not contain constant terms and linear terms.  $\square$

**Theorem 10.2.** Let  $T: \mathcal{P}_m \rightarrow \mathcal{P}_m$  be a map satisfying (7.8) and (10.1). Then  $r_n(\mu, T\mu)$  satisfy the conditions (C1) and (C2') w.r.t. the convolution  $\triangleright_T$ .

*Proof.* (C2') follows from the definition of conditionally monotone cumulants and (10.1). (C1) is showed as follows:

$$r_n^T(\mu^{\triangleright_T N}) = r_n(\mu^{\triangleright_T N}, T(\mu^{\triangleright_T N})) = r_n((\mu, T\mu)^{\triangleright N}) = Nr_n(\mu, T\mu). \quad \square$$

We note that the conditionally free cumulants  $R_n(\mu, T\mu)$  satisfy the conditions (C1) and (C2') under the same restriction, but this fact will not be used in this paper.

**Proposition 10.3.** Let  $T: \mathcal{P}_m \rightarrow \mathcal{P}_m$  be a map satisfying the condition (7.4). We assume that the  $n$ -th moment of  $T\mu$  is of the form

$$m_n(T\mu) = V_n(m_1(\mu), \dots, m_{n+1}(\mu)) \quad (10.2)$$

for any  $n \geq 1$ , where  $V_n$  is a polynomial which does not contain a constant term. Then the convolution  $\boxplus_T$  satisfies the conditions (M1) and (M2), and  $R_n(\mu, T\mu)$  satisfy the conditions (C1) and (C2').

**Remark 10.4.** The condition (10.2) is satisfied in all convolutions studied in [7, 8, 15, 16, 28].

## 11 Limit theorems of deformed convolutions

Central limit theorem and Poisson's law of small numbers of the convolution  $\triangleright_{\tau}$  hold by Theorem 4.10 in the sense of the convergence of moments under the conditions (7.8) and (10.1). We summarize the statements combining Theorem 4.10 and Theorem 10.2.

**Theorem 11.1.** *Let  $T: \mathcal{P}_{\mathfrak{m}} \rightarrow \mathcal{P}_{\mathfrak{m}}$  be a map which satisfies (7.8) and (10.1).*

(1) (central limit theorem) *Let  $\mu$  be a probability measure in  $\mathcal{P}_{\mathfrak{m}}$  with mean 0 and variance 1. We define  $\mu_N := (D_{\frac{1}{\sqrt{N}}}\mu)^{\triangleright_{\tau} N}$ . Then  $m_n(\mu_N)$  converges to  $m_n(\nu_1^{(T)})$ , where  $m_n(\nu_t^{(T)})$  are characterized by*

$$\frac{\partial}{\partial t} H_{\nu_t^{(T)}}(z) = -\frac{1}{H_{T\nu_t^{(T)}}(z)}. \quad (11.1)$$

(2) (Poisson's law of small numbers) *Let  $\{\mu^{(N)}\}_{N=1}^{\infty}$  be a sequence of probability measures in  $\mathcal{P}_{\mathfrak{m}}$  such that  $Nm_n(\mu^{(N)}) \rightarrow \lambda > 0$  as  $N \rightarrow \infty$  for all  $n \geq 1$ . We define  $\mu_N := (\mu^{(N)})^{\triangleright_{\tau} N}$ . Then  $m_n(\mu_N)$  converges to  $m_n(p_{\lambda}^{(T)})$ , where  $m_n(p_{\lambda}^{(T)})$  are characterized by*

$$\frac{\partial}{\partial \lambda} H_{p_{\lambda}^{(T)}}(z) = \frac{H_{Tp_{\lambda}^{(T)}}(z)}{1 - H_{Tp_{\lambda}^{(T)}}(z)}. \quad (11.2)$$

In this section we calculate the limit distributions for  $T$  constructed in the sections 8 and 9. If  $T$  is invertible, we can use monotone cumulants to calculate the limit distributions since  $\mu \triangleright_{\tau} \nu = T^{-1}(T\mu \triangleright T\nu)$ . Cumulants introduced in Section 10, however, enable us to calculate the limit distributions for even non-invertible  $T$ . In this section, we always use cumulants introduced in Section 10.

### 11.1 $V_{t,u,a}$ -transformation

We now calculate the limit distribution in central limit theorem of the convolution  $\triangleright_{t,u,a}$  for  $t > 0$ ,  $u \in \mathbb{R}$  and  $a \in \mathbb{R}$ . We only calculate the cases  $a = 0$  and  $t = 0$ . We denote  $r_n^{V_{t,u,a}}(\mu)$  by  $r_n^{(t,u,a)}(\mu)$ . In this section we use two logarithms  $\log_{[1]}$  and  $\log_{[2]}$ :  $\log_{[1]}(z)$  is defined by  $\log_{[1]}(z) := \log|z| + i \arg(z)$ ,  $\arg(z) \in (-\pi, \pi)$ ,  $z \in \mathbb{C} \setminus (-\infty, 0]$ ;  $\log_{[2]}$  is defined by  $\log_{[2]}(z) := \log|z| + i \arg(z)$ ,  $\arg(z) \in (0, 2\pi)$ ,  $z \in \mathbb{C} \setminus [0, \infty)$ . We define  $\sqrt{z}$  by  $\exp(\frac{1}{2} \log_{[2]} z)$  for  $z \in \mathbb{C} \setminus [0, \infty)$ . Then the Cauchy transform of the normalized arcsine law is given by  $\frac{1}{\sqrt{z^2 - 2}}$  for  $z \in \mathbb{C}_+$ .

**Theorem 11.2.** (1) *Let  $\mu$  be a probability measure in  $\mathcal{P}_{\mathfrak{m}}$  with mean 0 and variance 1. Then  $\mu_N := (D_{\frac{1}{\sqrt{N}}}\mu)^{\triangleright_{t,u,0} N}$  converges weakly to the Kesten distribution  $\nu^{(t,0)}$ . For  $t \geq 1$ , the absolutely continuous part is  $\frac{1}{2\pi} \frac{\sqrt{2t-x^2}}{1-(1-\frac{t}{2})x^2} dx$  and is supported on  $[-\sqrt{2t}, \sqrt{2t}]$ . For  $t < 1$ ,  $\nu^{(t,0)}$  contains atoms at  $x = \pm \frac{1}{\sqrt{1-\frac{t}{2}}}$ .*  
(2) *Let  $\mu$  be a probability measure in  $\mathcal{P}_{\mathfrak{m}}$  with mean 0 and variance 1. Then  $\mu_N := (D_{\frac{1}{\sqrt{N}}}\mu)^{\triangleright_{0,0,a} N}$  converges weakly to a probability measure  $\nu^{(0,a)}$ . The absolutely continuous part of  $\nu^{(0,a)}$  is given by*

$$\nu^{(0,a)}|_{ac} = \begin{cases} \frac{a}{(\log|1+\frac{a}{x}| - ax)^2 + \pi^2} dx, & x \in [-a, 0], \quad a > 0, \\ \frac{|a|}{(\log|1+\frac{a}{x}| - ax)^2 + \pi^2} dx, & x \in [0, |a|], \quad a < 0. \end{cases} \quad (11.3)$$

$\nu^{(0,a)}$  contains two atoms: one in  $(-\infty, -a)$  and one in  $(0, \infty)$  for  $a > 0$ ; one in  $(-\infty, 0)$  and one in  $(|a|, \infty)$  for  $a < 0$ .

**Remark 11.3.** (1) The Kesten distribution also appears in the central limit theorem of  $\boxplus_{\mathcal{U}_t}$  [7, 8] with the parameter  $t$  replaced by  $2t$ .

(2) The limit distribution of (2) is symmetric only in the case of  $a = 0$  where the convolution is Boolean convolution (cf. Proposition 8.5).

*Proof.* (1) We denote  $r_n^{V_{t,u,a}}(\nu)$  by  $r_n^{(t,u,a)}(\nu)$ . Let  $\{\nu_s^{(t,0)}\}_{s \geq 0}$  be a (formal) convolution semigroup which is a solution of (11.1) for  $T = V_{t,u,0}$ . (The word “formal” means that the limit moments might not



be deterministic. Therefore, we consider  $\nu_s^{(t,0)}$  as a sequence of moments.) We note that  $m_1(\nu_s^{(t,0)}) = sr_1^{(t,u,0)}(\nu_1^{(t,0)}) = 0$ . Then  $H_{V_{t,u,a}\nu_s^{(t,a)}}(z) = tH_{\nu_s^{(t,a)}}(z) + (1-t)z$ . We denote  $H_{\nu_s^{(t,0)}}(z)$  by  $H_s(z)$ . (11.1) can be integrated and we obtain  $\frac{t}{2}H_s(z)^2 + (1-t)zH_s(z) = -s + (1-\frac{t}{2})z^2$ , which implies

$$G_s(z) = \frac{(\frac{1}{2} - \frac{t}{2}) + \frac{1}{2}\sqrt{z^2 - 2st}}{(1 - \frac{t}{2})z^2 - 2s}. \quad (11.4)$$

$G_1$  is the Cauchy transform of the Kesten distribution (see [8]), whose support is compact. Then the weak convergence holds (see Theorem 4.5.5 in [9]).

(2) Let  $\{\nu_s^{(0,a)}\}_{s \geq 0}$  be a (formal) convolution semigroup which is a solution of (11.1) for  $T = V_{0,0,a}$ . We note that  $m_1(\nu_s^{(0,a)}) = 0$  and  $\sigma^2(\nu_s^{(0,a)}) = r_2^{(0,0,a)}(\nu_s^{(0,a)}) = s$ . Then  $H_{V_{0,0,a}\nu_s^{(0,a)}}(z) = z + as$ . We denote  $H_{\nu_s^{(0,a)}}(z)$  by  $H_s(z)$  for simplicity. We have

$$\begin{aligned} H_s(z) &= - \int_0^s \frac{1}{z + ar} dr + z \\ &= z - \frac{1}{a} \log_{[1]} \left( 1 + \frac{as}{z} \right). \end{aligned} \quad (11.5)$$

**case  $a > 0$ :** The absolutely continuous part of the limit distribution is given by  $\frac{a}{(\log |1 + \frac{a}{x}| - ax)^2 + \pi^2} dx$  which is supported on the set  $\{x \in \mathbb{R}; G_1(x + i0) < 0\} = [-a, 0]$ . We can show that the limit distribution contains one atom in  $(0, \infty)$  and one atom in  $(-\infty, -a)$ .

**case  $a < 0$ :** The absolutely continuous part is given by  $\frac{|a|}{(\log |1 + \frac{a}{x}| - ax)^2 + \pi^2} dx$  which is supported on the set  $[0, |a|]$ . We can show that the limit distribution contains one atom in  $(|a|, \infty)$  and one atom in  $(-\infty, 0)$ .

We note that the case  $a = 0$  corresponds to the Boolean convolution, and hence, the limit distribution is  $\frac{1}{2}(\delta_{-1} + \delta_1)$ .  $\square$

We calculate the limit distribution of Poisson's law of small numbers. We consider only the case  $T = V_{0,u,a}$ ; otherwise, the explicit form is difficult to obtain.

**Theorem 11.4.** *Let  $\{\mu^{(N)}\}_{N=1}^\infty$  be a sequence of probability measures in  $\mathcal{P}_m$  such that  $Nm_n(\mu^{(N)}) \rightarrow \lambda > 0$  as  $N \rightarrow \infty$  for all  $n \geq 1$ . Then  $\mu_N := (\mu^{(N)})^{\triangleright_{0,u,a}N}$  converges weakly to a compactly supported distribution  $p_\lambda^{(u,a)}$ . The absolutely continuous part of  $p_\lambda^{(u,a)}$  is given by*

$$p_\lambda^{(u,a)}|_{ac} = \begin{cases} \frac{a-u}{\left(\log |1 + \frac{(a-u)\lambda}{x-1}| - (a-u)(x-\lambda)\right)^2 + \pi^2} dx, & x \in [1 - (a-u)\lambda, 1], \quad a > u, \\ \frac{|a-u|}{\left(\log |1 + \frac{(a-u)\lambda}{x-1}| - (a-u)(x-\lambda)\right)^2 + \pi^2} dx, & x \in [1, 1 + (u-a)\lambda], \quad a < u. \end{cases} \quad (11.6)$$

$p_\lambda^{(u,a)}$  contains two atoms: one in  $(-\infty, 1 - (a-u)\lambda)$  and one in  $(1, \infty)$  for  $a > u$ ; one in  $(-\infty, 1)$  and one in  $(1 + (u-a)\lambda, \infty)$  for  $a < u$ .

**Remark 11.5.** One can see that  $p_\lambda^{(u,a)}$  is in  $\mathcal{P}_+$  if and only if  $u \geq a$  (cf. Proposition 8.5).

*Proof.* We note that  $m_1(p_\lambda^{(u,a)}) = r_1^{(0,u,a)}(p_\lambda^{(u,a)}) = \lambda$  and  $\sigma^2(p_\lambda^{(u,a)}) = \lambda$ . Then we obtain the differential equation

$$\frac{\partial}{\partial \lambda} H_{p_\lambda^{(u,a)}}(z) = -1 - \frac{1}{z - 1 + (a-u)\lambda}. \quad (11.7)$$

The remaining arguments are similar to Theorem 11.2 and we omit the proof.  $\square$

## 11.2 Deformations related to $\triangleright$ -infinitely divisible distributions

For a compactly supported  $\triangleright$ -infinitely divisible distribution  $\xi$ , let  $\{\xi_t\}_{t \geq 0}$  be the corresponding weakly continuous  $\triangleright$ -convolution semigroup with  $\xi_0 = \delta_0$  and  $\xi_1 = \xi$ . Then  $\xi_t$  is compactly supported for each  $t > 0$  [24]. Let  $\{\nu_s^{[\xi,t]}\}_{s \geq 0}$  and  $\{p_s^{[\xi,t]}\}_{s \geq 0}$  be the (formal) convolution semigroups defined by (11.1)

and (11.2), respectively. We denote  $r_n^{\Xi_t^\xi}(\mu)$  by  $r_n^{[\xi, t]}(\mu)$ . Since  $r_2^{[\xi, t]}(\nu_s^{[\xi, t]}) = \sigma^2(\nu_s^{[\xi, t]}) = s$  we obtain  $\Xi_t^\xi(\nu_s^{[\xi, t]}) = \xi_{st}$ . Similarly, we obtain  $\Xi_t^\xi(p_\lambda^{[\xi, t]}) = \xi_{\lambda t}$ . Therefore, (11.1) and (11.2) become

$$\frac{\partial}{\partial s} H_{\nu_s^{[\xi, t]}}(z) = -\frac{1}{H_{\xi_{st}}(z)}, \quad (11.8)$$

$$\frac{\partial}{\partial \lambda} H_{p_\lambda^{[\xi, t]}}(z) = \frac{H_{\xi_{\lambda t}}(z)}{1 - H_{\xi_{\lambda t}}(z)}. \quad (11.9)$$

These equations have been defined in the sense of formal power series of moments. Moreover, in the present case, we can show that any one of (11.8) and (11.9) gives a sequence of moments of a compactly supported probability measure for each  $s > 0$  since the right hand sides of (11.8) and (11.9) are analytic outside a ball. Therefore,  $\nu_s^{[\xi, t]}$  and  $p_\lambda^{[\xi, t]}$  make sense as uniquely determined probability measures. Since the limit distribution is compactly supported, the convergence of moments in Theorem 11.1 becomes the weak convergence. We summarize the above arguments. We denote  $\triangleright_{\Xi_t^\xi}$  by  $\triangleright_{[\xi, t]}$ .

**Theorem 11.6.** *Let  $\xi$  be a  $\triangleright$ -infinitely divisible distribution in  $\mathcal{P}_c$ .*

(1) (central limit theorem) *Let  $\mu$  be a probability measure in  $\mathcal{P}_m$  with mean 0 and variance 1. Then  $\mu_N := (D_{\frac{1}{\sqrt{N}}} \mu)^{\triangleright_{[\xi, t]}^N}$  converges to  $\nu_1^{[\xi, t]}$  weakly.*

(2) (Poisson's law of small numbers) *Let  $\{\mu^{(N)}\}_{N=1}^\infty$  be a sequence of probability measures in  $\mathcal{P}_m$  such that  $Nm_n(\mu^{(N)}) \rightarrow \lambda > 0$  as  $N \rightarrow \infty$  for all  $n \geq 1$ . Then  $\mu_N := (\mu^{(N)})^{\triangleright_{[\xi, t]}^N}$  converges to  $p_\lambda^{[\xi, t]}$  weakly.*

We calculate the limit distributions explicitly when  $\xi$  is the normalized arcsine law.

**Theorem 11.7.** *Let  $\eta$  be the normalized arcsine law.*

(1) *The limit distribution  $\nu_1^{[\eta, t]}$  is given by the Kesten distribution  $\nu^{(t, 0)}$ .*  
(2) *The absolutely continuous part of  $p_\lambda^{[\eta, t]}$  is supported on  $[-\sqrt{2\lambda t}, \sqrt{2\lambda t}] \cup [1, \sqrt{2\lambda t + 1}]$ . The singular part consists of three parts: one atom exists in  $(\sqrt{2\lambda t + 1}, \infty)$ ; one atom exists in  $(\sqrt{2\lambda t}, 1)$  if  $0 < t < \frac{1}{2\lambda}$  and  $(1 - \frac{1}{t})\sqrt{2\lambda t} - \frac{1}{t} \log(1 - \sqrt{2\lambda t}) - \lambda < 0$ ; one atom exists in  $(-\infty, -\sqrt{2\lambda t})$  if  $(\frac{1}{t} - 1)\sqrt{2\lambda t} - \frac{1}{t} \log(\sqrt{2\lambda t + 1}) > 0$ .*

**Remark 11.8.** It is remarkable that the limit distribution in (1) also appears in Theorem 10 in [15] with the parameter  $t$  replaced by  $2t$ .

*Proof.* (1) The differential equation (11.8) becomes

$$\frac{\partial}{\partial s} H_{\nu_s^{[\eta, t]}}(z) = -\frac{1}{\sqrt{z^2 - 2ts}}, \quad (11.10)$$

which implies  $H_{\nu_s^{[\eta, t]}}(z) = (1 - \frac{1}{t})z + \frac{1}{t}\sqrt{z^2 - 2ts}$ . Therefore, the limit distribution is the Kesten distribution.

(2) The differential equation (11.9) becomes

$$\frac{\partial}{\partial \lambda} H_{p_\lambda^{[\xi, t]}}(z) = \frac{\sqrt{z^2 - 2\lambda t}}{1 - \sqrt{z^2 - 2\lambda t}}. \quad (11.11)$$

We can solve this and obtain

$$H_{p_\lambda^{[\xi, t]}}(z) = \left(1 - \frac{1}{t}\right)z + \frac{1}{t}\sqrt{z^2 - 2\lambda t} + \frac{1}{t} \log_{[1]} \left( \frac{\sqrt{z^2 - 2\lambda t} - 1}{z - 1} \right) - \lambda. \quad (11.12)$$

One can see that  $\lim_{\text{Im } z \searrow 0} H_{p_\lambda^{[\xi, t]}}(z) > 0$  if and only if  $\text{Re } z \in (-\sqrt{2\lambda t}, \sqrt{2\lambda t}) \cup [1, \sqrt{2\lambda t + 1}]$ . We remark that  $H_{p_\lambda^{[\xi, t]}}(x)$  is strictly increasing in the intervals  $(-\infty, -\sqrt{2\lambda t})$ ,  $(\sqrt{2\lambda t}, 1)$  and  $(\sqrt{2\lambda t + 1}, \infty)$ . Then it is not difficult to show the existence of atoms.  $\square$

# Acknowledgements

The author would like to thank Prof. Marek Bożejko for guiding him to the notion of conditionally free independence and an important reference [13]. He thanks Prof. Uwe Franz for fruitful discussions and for giving a seminar on the categorical treatment of independence during the visit to Kyoto. He is also grateful to Prof. Izumi Ojima, Prof. Shogo Tanimura, Mr. Hayato Saigo, Mr. Ryo Harada, Mr. Hiroshi Ando and Mr. Kazuya Okamura for their comments and encouragements. This work was supported by JSPS KAKENHI 21-5106.

# References

- [1] L. Accardi, R. Lenczewski and R. Śaławapata, Decompositions of the free product of graphs, *Infin. Dim. Anal. Quantum Probab. Rel. Topics*, **10**, no. 3 (2007), 303-334.
- [2] O. E. Barndorff-Nielsen, U. Franz, R. Gohm, B. Kümmerer and S. Thorbjørnsen, *Quantum Independent Increment Processes II*, Lecture Notes in Math., vol. 1866, Springer-Verlag, 2006.
- [3] A. Ben Ghorbal and M. Schürmann, Non-commutative notions of stochastic independence, *Math. Proc. Comb. Phil. Soc.* **133** (2002), 531-561.
- [4] H. Bercovici, Multiplicative monotonic convolution, *Illinois J. Math.* **49**, no.3 (2005), 929-951.
- [5] H. Bercovici, On Boolean convolutions, *Operator Theory* **20**, 7-13, Theta. Ser. Adv. Math. 6, Theta, Bucharest, 2006.
- [6] M. Bożejko, M. Leinert and R. Speicher, Convolution and limit theorems for conditionally free random variables, *Pac. J. Math.* **175** (1996), no. 2, 357-388.
- [7] M. Bożejko and J. Wysoczański, New examples of convolutions and non-commutative central limit theorems, *Banach Center Publ.*, **43** (1998), 95-103.
- [8] M. Bożejko and J. Wysoczański, Remarks on  $t$ -transformations of measures and convolutions, *Ann. I. H. Poincaré-PR* **37** (2001), 737-761.
- [9] K. L. Chung, *A Course in Probability Theory*, Harcourt, Brace & World, Inc., 1968.
- [10] T. Hasebe, Monotone convolution semigroup, in preparation.
- [11] T. Hasebe and H. Saigo, The monotone cumulants, available in arXiv:0907.4896v1.
- [12] H. Kesten, Symmetric random walks on groups, *Trans. Amer. Math. Soc.* **92**, no. 2 (1959), 336-354.
- [13] U. Franz, Multiplicative monotone convolutions, *Banach Center Publ.*, **73** (2006), 153-166.
- [14] U. Franz, Boolean convolution of probability measures on the unit circle, available in arxiv:math/0403243v2.
- [15] A. D. Krystek and Ł. J. Wojakowski, Associative convolutions arising from conditionally free convolution, *Inf. Dim. Anal. Quantum Probab. Rel. Topics* **8**, no. 3 (2005), 515-545.
- [16] A. D. Krystek and H. Yoshida, Generalized  $t$ -transformations of probability measures and deformed convolution, *Probab. Math. Stat.* **24** (2004), 97-119.
- [17] F. Lehner, Cumulants in noncommutative probability theory I, *Math. Z.* **248** (2004), 67-100.
- [18] R. Lenczewski and R. Śaławapata, Discrete interpolation between monotone probability and free probability, *Infin. Dim. Anal. Quantum Probab. Rel. Topics*, **9**, no. 1 (2006), 77-106.
- [19] R. Lenczewski, Decompositions of the additive free convolution, *J. Funct. Anal.* **246** (2007), 330-365.

- [20] R. Lenczewski, Operators related to subordination for free multiplicative convolutions, *Indiana Univ. Math. J.* **57**, no. 3 (2008), 1055-1103.
- [21] R. Lenczewski and R. Śaławata, Discrete interpolation between monotone probability and free probability, *Infin. Dim. Anal. Quantum Probab. Rel. Topics*, **9**, no. 1 (2006), 77-106.
- [22] R. Lenczewski and R. Śaławata, Noncommutative Brownian motions associated with Kesten distributions and related Poisson processes, *Infin. Dim. Anal. Quantum Probab. Rel. Topics*, **11**, no. 3 (2008), 351-375.
- [23] H. Maassen, Addition of freely independent random variables, *J. Funct. Anal.* **106** (1992), 409-438.
- [24] N. Muraki, Monotonic convolution and monotonic Lévy-Hinčin formula, preprint, 2000.
- [25] N. Muraki, The five independences as quasi-universal products, *Infin. Dim. Anal. Quantum Probab. Rel. Topics*, **5**, no. 1 (2002), 113-134.
- [26] N. Muraki, The five independences as natural products, *Infin. Dim. Anal. Quantum Probab. Rel. Topics*, **6**, no. 3 (2003), 337-371.
- [27] A. Nica and R. Speicher, *Lectures on the Combinatorics of Free Probability*, London Math. Soc. Lecture Note Series, vol. 335, Cambridge Univ. Press, 2006.
- [28] F. Oravecz, Fermi convolution, *Infin. Dim. Anal. Quantum Probab. Rel. Topics*, **5**, no. 2 (2002), 235-242.
- [29] F. Oravecz, The number of pure convolutions arising from conditionally free convolution, *Infin. Dim. Anal. Quantum Probab. Rel. Topics*, **8**, no. 3 (2005), 327-355.
- [30] A. N. Shiryaev, *Probability*, Springer-Verlag, New York, 1984.
- [31] R. Speicher, On universal products, in *Free Probability Theory*, papers from a Workshop on Random Matrices and Operator Algebra Free Products, Toronto, Canada 1995, ed. D. V. Voiculescu, Fields Inst. Commun. 12 (Amer. Math. Soc., 1997), 257-266.
- [32] R. Speicher and R. Woroudi, Boolean convolution, in *Free Probability Theory*, papers from a Workshop on Random Matrices and Operator Algebra Free Products, Toronto, Canada 1995, ed. D. V. Voiculescu, Fields Inst. Commun. 12 (Amer. Math. Soc., 1997), 267-280.
- [33] D. Voiculescu, K. J. Dykema and A. Nica, *Free Random Variables*, CRM Monograph Series, AMS, 1992.